

La découverte d'une nouvelle espèce de la série *guyanensis*, en sympatrie avec *P. geniculatus* et *P. corossoniensis* dans le bassin du rio Napo, est en accord avec l'écologie de ces espèces qui affectionnent la forêt primaire très arrosée ; en Guyane française, également on trouve ensemble ces deux mêmes espèces et *P. dorlinsis*, cela risquant d'ailleurs de compliquer la tâche pour qui s'intéressera à la réhabilitation de *P. guyanensis* Floch & Abonnenc.

REMERCIEMENTS. – Nous remercions Madame A. Aing de l'IRD, Bondy (Institut de Recherche d'Ile-de-France) pour avoir effectué la mise en page des illustrations.

AUTEURS CITÉS

- ABONNENC E., 1972. – Les phlébotomes de la région éthiopienne (Diptera, Psychodidae). *Mémoires ORSTOM*, **55**, 289 p.
- AMUNNARIZ M., 1984. – *Salud Y Enfermedad. Patologia Tropical en la Region Amazonica Ecuatoriana*. CICAME, 98 p.
- GALATI E. A. B., 1995. – Phylogenetic systematics of Phlebotominae (Diptera, Psychodidae) with emphasis on American groups. Proceedings of the II International Symposium on Phlebotomine Sandflies. *Boletin de la Direccion de Malariologia y Saneamiento Ambiental*, **35** (Suppl. 1) : 133-142.
- YOUNG D. G. & ROGERS T. E., 1984. – The phlebotomine sand fly fauna (Diptera: Psychodidae) of Ecuador. *Journal of Medical Entomology*, **21** : 597-611.

Yair BEN-DOV & Emilio GUERRIERI. – *Waxiella mimosae* (Signoret) (Hemiptera, Coccoidea, Coccidae) and its parasitoid *Anicetus africanus* (Girault) (Hymenoptera, Encyrtidae) newly recorded from Israel

Wax scale insects, comprising about 150 species and subspecies, are grouped in the subfamily Ceroplastinae in the soft scale insect family Coccidae (DE LOTTO, 1965; GIMPEL *et al.*, 1974; BEN-DOV, 1986). Most species of wax scales were placed in the senior genus *Ceroplastes* Gray, while some additional genera were introduced, namely, *Gascardia* Targioni Tozzetti, *Vinsonia* Signoret, and *Waxiella* De Lotto. The 20 species and subspecies, which have been assigned to the genus *Waxiella*, are distributed mainly in the Afrotropical region (DE LOTTO, 1969, 1971; BEN-DOV, 1993, 2008).

In this note we record for the first time in Israel, the wax scale *Waxiella mimosae* (Signoret) and its parasitoid *Anicetus africanus* (Girault). *Waxiella mimosae*, which was originally described from Egypt, has been recorded also from widely-separated territories all in the Afrotropical region, namely, South Africa (ANNECKE, 1967), Angola (ALMEIDA, 1973), Mauritania (BALACHOWSKY & MATILE-FERRERO, 1971), Algeria (BALACHOWSKY, 1934), and Saudi Arabia (MATILE-FERRERO, 1988).

Waxiella mimosae is found so far in Israel at the southern Arava Valley, which is part of the Great Rift Valley. The Arava Valley stretches from Eilat city at the south to the Dead Sea in the north. The climate is very arid with an average annual rainfall of 20 millimetres and with large annual variations in temperature. The mean maximal temperature fluctuates between 39.9°C in July and 20.8°C in January. The average annual relative humidity ranges from 15% (in June) to 33 % in December. We record *W. mimosae* from two species of *Acacia*, as well as on the green semiparasitic mistletoe *Loranthus acaciae* (Loranthaceae), which is a common parasite of *Acacia* trees in this region.

Waxiella mimosae has two annual generations in the Arava valley. No males were observed, and we assume that this species reproduces parthenogenetically. Adult females of the first generation appear in March, and reproduction of the second generation takes place in October.

Anicetus africanus has been reported from *W. mimosae* in South Africa by ANNECKE (1967), who suggested that the earlier record by MERCET (1925) of this parasitoid from *Ceroplastes africana* in Egypt needs further confirmation. *Anicetus africanus* is a distinctive parasitoid whose females can be identified by a combination of characters as reported by ANNECKE (1967): general colour orange weakly metallic on thorax and antenna, these latter strongly expanded and flattened with pedicel overlapping the first funicular segment; ovipositor strongly exerted, the exerted part 0.5x as long as the gaster. The rate of parasitization in a sample collected on 19.x.2008 has reached 30 % with a balanced sex ratio (1:1). In about 80% of the not-parasitized females, encapsulation bodies were found.

Material examined: *Waxiella mimosae*: all specimens collected in Israel, and deposited in the Coccoidea Collection, Department of Entomology, Bet Dagan, Israel. Yotvata, 10.iii.2008, *Acacia tortilis* (Fabaceae) (C-4428); Yotvata, 10.iii.2008, *Loranthus acaciae* (Loranthaceae) (C-4429); Yotvata, 10.vi.2008, *L. acaciae* (C-4474); Yotvata, 22.vii.2008, *L. acaciae* (C-4489); Nahal Qetura, 22.iv.1986, *A. raddiana* (C-4494); Yotvata, 19.x.2008, *L. acaciae* (C-4539). *Anicetus africanus*: Israel, Yotvata 19.x.2008 ex *Waxiella mimosae* on *Loranthus acaciae* parasitizing *Acacia tortilis* (Y. Ben Dov). Specimens of *A. africanus* are deposited in the collection of Hymenoptera at the Dipartimento di Entomologia e Zoologia Agraria Filippo Silvestri”, Università degli Studi di Napoli “Federico II” Portici, Naples, Italy and at The Natural History Museum, London, UK.

REFERENCES

- ALMEIDA D. M. DE, 1973. – Coccoidea de Angola. 1. Revisão das especies conhecidas. *Boletim do Instituto de Investigação Científica de Angola*, **10**: 1-23.
- ANNECKE D. P., 1967. – The genera *Anicetus* Howard, 1896, *Paracerapterocerus* Girault, 1920, and allies, with descriptions of new genera and species (Hymenoptera: Encyrtidae). *Transactions of the Royal Entomological Society of London*, **119**: 99-169.
- BALACHOWSKY A. S., 1934. – Les coccides du Sahara central. Mission du Hoggar. III. (Février à mai 1928). In: Seurat L.-G., Etudes zoologiques sur le Sahara central. *Mémoires de la Société d'Histoire Naturelle de l'Afrique du Nord*, **4**: 145-157.
- BALACHOWSKY A. S. & MATILE-FERRERO D., 1971. – Les Cochenilles (Hom., Coccoidea) de la République Islamique de Mauritanie. Contribution à l'étude des Coccoidea de l'Afrique tropicale et équatoriale (14^e note). *Bulletin de l'Institut fondamental d'Afrique noire, série A*, **32** (4) : 1078-1087.
- BEN-DOV Y., 1986. – Taxonomy of two described and one new species of *Waxiella* De Lotto (Homoptera: Coccoidea: Coccidae). *Systematic Entomology*, **11**: 165-174.
- 1993. – *A systematic catalogue of the soft scale insects of the world (Homoptera: Coccoidea: Coccidae) with data on geographical distribution, host plants, biology and economic importance*. Gainesville, Florida: Sandhill Crane Press, 536 p.
- 2008. – ScaleNet, Family Coccidae. December, 2008. www.sel.barc.usda.gov/catalogs/Coccidae.htm
- DE LOTTO G., 1965. – On some Coccidae (Homoptera), chiefly from Africa. *Bulletin of the British Museum (Natural History), Entomology*, **16**: 175-239.
- 1969. – A new genus of wax scales (Homoptera: Coccidae). *Bolletino del Laboratorio di Entomologia Agraria "Filippo Silvestri" di Portici*, **27**: 210-218.
- 1971. – On some genera and species of wax scales (Homoptera: Coccidae). *Journal of Natural History*, **5**: 133-153.
- GIMPEL W. F., MILLER D.R. & DAVIDSON J. A., 1974. – A systematic revision of the wax scales, genus *Ceroplastes*, in the United States (Homoptera: Coccoidea: Coccidae). *University of Maryland, Agricultural Experiment Station, Miscellaneous Publication*, **841**: 1-85.
- MATILE-FERRERO D., 1988. – Sternorrhyncha: Suborder Coccoidea of Saudi Arabia (Part 2). *Fauna of Saudi Arabia*, **9**: 23-38.
- MERCET R. G., 1925. – Encyrtides et Aphelinides d'Egypte. *Bulletin de la Société Royale Entomologique d'Egypte*, **1925**: 46-55.
- (Y. B-D.: Department of Entomology, Agricultural Research Organization, The Volcani Center, Bet Dagan 50250, Israël <yairbd@netvision.net.il> ; E. G.: Institute for Plant Protection, National Research Council of Italy, Via Università, 133, 80055 – Portici (NA), Italy <guerrieri@ipp.cnr.it>)