

## New hover fly records for Corsica: results from *Our Planet Reviewed in Corsica 2019-2021* (Diptera, Syrphidae)

Ximo MENGUAL<sup>1</sup>, Thomas LEBARD<sup>2</sup> & Alexandre CORNUEL-WILLERMOZ<sup>3</sup>

<sup>1</sup> Museum Koenig, Leibniz-Institut zur Analyse des Biodiversitätswandels, Adenauerallee 127,  
D – 53113 Bonn, Germany <x.mengual@leibniz-lib.de>

<sup>2</sup> Quartier Ginestrea, hameau de Piène-Haute, F – 06540 Breil-sur-Roya, France <thomas\_lebard@yahoo.fr>

<sup>3</sup> Office de l’Environnement de la Corse – Observatoire Conservatoire des Invertébrés de Corse,  
14 avenue Jean-Nicoli, F – 20250 Corte, France <alexandre.cornuel-willermoz@oec.fr>

(Accepté le 2.XI.2023 ; publié en ligne le 15.XII.2023)

**Abstract.** – In this study we present a commented and referenced species list of the Syrphidae (Diptera) collected during the surveys of the *Our Planet Reviewed in Corsica 2019-2021* programme. We studied a total of 2420 specimens belonging to 131 taxa, which were collected during the field campaigns between 2019 and 2022 using different methodologies: hand collecting, hand-net, interception traps, light traps, Malaise traps, sweep-netting, and coloured pan traps. The genera *Anasimyia* Schiner, 1864, *Brachyopa* Meigen, 1822, *Lapposyrphus* Dušek & Láska, 1967, *Lejops* Rondani, 1857, *Melangyna* Verrall, 1901, *Neocnemodon* Goffe, 1944, and *Parhelophilus* Girschner, 1897, are recorded from Corsica for the first time, and we also provide records for 30 species new to Corsica. Among the studied material, we present records for 17 threatened species in Europe and three endemics of Corsica. Furthermore, we report a new location for the Critically Endangered *Riponnensis daccordii* (Claussen, 1991) and corroborate that it is not extinct.

**Résumé.** – Nouvelles données sur les syrphes pour la Corse : résultats de *La Planète Revisée en Corse 2019-2021* (Diptera, Syrphidae). Nous présentons dans cet article une liste commentée et référencée des espèces de Syrphidae (Diptères) collectées lors des suivis du programme *La Planète Revisée en Corse 2019-2021*. Nous avons étudié un total de 2420 spécimens appartenant à 131 taxons, qui ont été collectés lors des campagnes de terrain entre 2019 et 2022 en utilisant différentes méthodologies : assiettes colorées, collecte manuelle, filet à main, filet fauchoir, pièges d’interception, pièges lumineux et pièges Malaise. Les genres *Anasimyia* Schiner, 1864, *Brachyopa* Meigen, 1822, *Lapposyrphus* Dušek & Láska, 1967, *Lejops* Rondani, 1857, *Melangyna* Verrall, 1901, *Neocnemodon* Goffe, 1944, et *Parhelophilus* Girschner, 1897, sont mentionnés en Corse pour la première fois, et nous fournissons également des données pour 30 espèces nouvelles pour la Corse. Parmi le matériel étudié, nous présentons les données de 17 espèces menacées en Europe et trois endémiques de Corse. En outre, nous signalons une nouvelle station pour *Riponnensis daccordii* (Claussen, 1991), espèce en danger critique d’extinction, et corroborons qu’elle n’est pas éteinte.

**Keywords.** – Flower flies, hoverflies, new species records, *Riponnensis daccordii*, endemic, threatened species.

Known as hover flies or flower flies, member of the family Syrphidae Latreille, 1802 (Insecta, Diptera) are popular among nature-lovers and entomologists. The family comprises over 6300 described species worldwide (SKEVINGTON *et al.*, 2019), with more than 900 species occurring in Europe (VUJIĆ *et al.*, 2022; STÅHLS, 2022) — a number that grows every year (e.g., BOT *et al.*, 2022; ARACIL *et al.*, 2023). The family is divided into four subfamilies (MENGUAL *et al.*, 2015a), namely Eristalinae Newman, 1834, Microdontinae Rondani, 1845, Pipizinae Williston, 1885, and Syrphinae Latreille, 1802. Recent phylogenetic studies find support for the monophyly of these subfamilies, except for Eristalinae (YOUNG *et al.*, 2016; PAULI *et al.*, 2018; MORAN *et al.*, 2022; MENGUAL *et al.*, 2023).

Syrphids are known for being excellent mimics of aculeate hymenopterans (ROTHRAY & GILBERT, 2011), but also very important for their ecosystem services (DUNN *et al.*, 2020). Adults feed on pollen and nectar and are important pollinators in natural and agricultural ecosystems (SSYMANEK & KEARNS, 2009; INOUYE *et al.*, 2015; DOYLE *et al.*, 2020). Hover fly

larvae show a broad array of feeding modes (ROTHRAY, 1993; ROTHERAY & GILBERT, 2011; PÉREZ-LACHAUD *et al.*, 2014; FLEISCHMANN *et al.*, 2016, 2022) and act as biological control agents of arthropod pests (ARCAYA *et al.*, 2017; BELLEFEUILLE *et al.*, 2019; MOERKENS *et al.*, 2021) and also of certain weeds (RIZZA *et al.*, 1988; SHEPPARD *et al.*, 1995; GROSSKOPF, 2005). Moreover, saprophagous larvae serve as decomposers of organic matter (LARDÉ, 1989, 1990; MORALES & WOLFF, 2010) and have a relative significance as insects of forensic importance (MAGNI *et al.*, 2013; HEO *et al.*, 2020) and as myiasis agents (PÉREZ-BAÑÓN *et al.*, 2020).

More than one third of the European hover flies are threatened with extinction (VUJIĆ *et al.*, 2022) and recent studies show a strong decline in their abundance (GATTER *et al.*, 2020; BARENDRUGT *et al.*, 2022). Now more than ever large-scale scientific studies are needed to understand the fluctuations and responses of the pollinator communities to environmental changes in Europe, and several European Commission-funded projects aim to provide such information, such as the EU Pollinator Monitoring Scheme (EUPoMS; see POTTS *et al.*, 2021), the Preparatory Action for EU Pollinator Monitoring Scheme and Indicators (SPRING project), the Horizon 2020 Europe research projects (PoshBee, Safeguard), and European National action plans for pollinators (see <https://wikis.ec.europa.eu/display/EUPKH/EU+Pollinator+Information+Hive>).

In the present study we report the new records of Syrphidae from Corsica resulting from the expeditions of the *Our Planet Reviewed in Corsica 2019–2021* programme (TOUROULT *et al.*, 2023). Although the discovery of new species records for Corsica was not among the main objectives of the programme, the amount of sampled material may provide a very accurate picture of the current fauna of this Mediterranean island and, hence, improve our knowledge on insects, their distribution and biology. Knowing that a large proportion of European hover flies are threatened and that Corsica houses several endemics, we believe that our study can help other European initiatives with pollinators and, wherever feasible, it will contribute to a better management of the territory through assessments and monitoring.

## MATERIAL AND METHODS

Specimens were collected in the frame of the most recent edition of the *La Planète Revisée* (*Our Planet Reviewed*) series of surveys (<https://www.mnhn.fr/fr/recherche-expertise/lieux/planete-revisitee>). Between 2019 and 2021, scientific field expeditions on the island of Corsica took place organised by the Muséum national d’Histoire naturelle (MNHN, Paris, France), the Office français de la biodiversité, and the Collectivité de Corse (see <http://laplaneterevisitee-corse.mnhn.fr>). A detailed protocol on the survey strategy, sampled localities, workflow and methodologies used during the field campaigns can be found in TOUROULT *et al.* (2023). In 2022, Thomas Lebard continued the field work within the frame of Our Planet Reviewed and visited Corsica for more sampling. Simplemappr (SHORTHOUSE, 2010) was used to create a map with the 267 sampling localities where hover flies were collected (fig. 1).

The studied and reported material can be divided into two parts or sections: those specimens collected by hand-net by Thomas Lebard, and all other samples originated from the field campaigns using several methodologies, such as hand-collecting, interception traps, light traps, Malaise traps, pan traps, and sweep-netting (see TOUROULT *et al.*, 2023). For the latter samples, the Diptera coordinator (Marc Pollet, Research Institute for Nature and Forest – INBO, Brussels, Belgium) sent the subsamples to the first author after sorting the specimens to family. The sampling methodology is indicated for each studied specimen (see Supplementary material). Species records will ultimately be disclosed and distributed in the frame of the Inventaire national du patrimoine naturel (<https://inpn.mnhn.fr/>) and, in the case of this specific




Fig. 1.—Map of Corsica with the sampling localities during the field campaigns of *La Planète Revisée (Our Planet Reviewed)*.

survey, via the collections portal of the MNHN ([https://science.mnhn.fr/institution/mnhn/item/search/form?lang=fr\\_FR](https://science.mnhn.fr/institution/mnhn/item/search/form?lang=fr_FR)).

For the morphological identification of the hover fly individuals, several literature works were used. For the genus identification we used the keys of VAN VEEN (2010) and SPEIGHT (2020a). For the identifications at the species level we also used VAN VEEN (2010) as a general reference, together with SPEIGHT & SARTHOU (2017) and the following additional references:

*Anasmyia* Schiner, 1864 (CLAUSSEN & TORP, 1980); *Brachyopa* Meigen, 1822 (PELLMANN, 1998); *Callicera* Panzer, 1809 (SPEIGHT, 1991; SMIT, 2014); *Ceriana* Rafinesque, 1815 (VAN STEENIS *et al.*, 2016); *Claussenia* Vujić & Ståhls, 2013 (VUJIĆ *et al.*, 2013); *Doros* Meigen, 1803 (SPEIGHT, 1988); *Eumerus* Meigen, 1822 (VAN DER GOOT, 1968; DOCZKAL, 1996; SPEIGHT *et al.*, 2021); *Merodon* Meigen, 1803 (VUJIĆ *et al.*, 2021; and references therein); *Myolepta* Newman, 1838 (REEMER *et al.*, 2004); *Neocnemodon* Goffe, 1944 (BARTSCH, 2009); *Paragus* Latreille, 1804 (GOELDLIN DE TIEFENAU, 1976; GOELDLIN DE TIEFENAU & LUCAS, 1981; MARCOS-GARCÍA, 1986; MARCOS-GARCÍA & ROJO, 1994; SOMMAGGIO, 2002); *Pelecocera* Meigen, 1822 (VAN ECK & MENGUAL, 2021; LAIR *et al.*, 2022); and *Riponnensis* Maibach, Goedlin de Tiefenau & Speight, 1994 (CLAUSSEN, 1991; BOT *et al.*, 2023).

## RESULTS

In this section we exclusively report the species collected in the frame of the programme *Our Planet Reviewed in Corsica 2019-2021* (TOUROULT *et al.*, 2023). Besides the hand-net, other methodologies that yielded specimens are hand collecting, interception traps (Polytrap™), light traps, Malaise traps, blue, yellow, pink and white pan traps, and sweep-netting.

For each collected species we provide the literature references for Corsica, the number of specimens divided by sex (see Supplementary material), known distribution (based on SPEIGHT, 2020b and IUCN, 2023), the IUCN status (after IUCN, 2023), and some remarks. For each literature reference or record, we cite the name of the taxon as published in the original publication, either if it is a synonym or an old combination. Studied specimens with

their metadata are listed in Supplementary material. A total of 2420 specimens belonging to 131 taxa were collected during the field campaigns between 2019 and 2022. Due to the preservation condition or the impossibility to distinguish the female sex among close species several specimens were left without species identification, namely individuals of the genera *Chrysotoxum* Meigen, 1803, *Eumerus*, *Myolepta*, *Neocnemodon*, *Paragus*, and *Platycheirus* Lepeletier de Saint-Fargeau & Audinet-Serville, 1828.

Authors were requested to organize the list by subfamilies, but the subfamilial division of this group will change soon (see MORAN *et al.*, 2022). Thus, the species are listed in alphabetic order.

***Anasimyia contracta*** Claussen & Torp, 1980

**Material examined.** – 3 males, 1 female.

**Distribution.** – British Isles, Europe, Turkey, and European parts of Russia.

**IUCN status.** – Least Concern.

**Remarks.** – New species and new genus for Corsica.

***Baccha elongata*** (Fabricius, 1775)

**References.** – BECKER *et al.* (1910); KUNTZE (1913) as *Baccha obscuripennis* Meigen, 1822; DIRICKX (1994) as *Baccha obscuripennis* and *B. elongata*; SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 21 males, 15 females.

**Distribution.** – British Isles, Europe, Azores, and European parts of Russia.

**IUCN status.** – Least Concern.

**Remarks.** – Based on our own research using morphological and molecular characters, we do consider *Baccha obscuripennis* a junior synonym of *B. elongata* following previous authors (SPEIGHT, 2020b; MENGUAL *et al.* (2020)).

***Brachyopa pilosa*** Collin, 1939

**Material examined.** – 1 male, 1 female.

**Distribution.** – Europe, Great Britain, and European parts of Russia.

**IUCN status.** – Least Concern.

**Remarks.** – New species and new genus for Corsica.

***Brachypalpooides lentus*** (Meigen, 1822)

**References.** – KUNTZE (1913) as *Xylota lenta*; DIRICKX (1994); SPEIGHT *et al.* (2020).

**Material examined.** – 1 male.

**Distribution.** – British Isles, Europe, Middle East, and European parts of Russia.

**IUCN status.** – Least Concern.

***Callicera aurata*** (Rossi, 1790)

**Material examined.** – 9 females.

**Distribution.** – British Isles, Europe, Turkey, southern parts of European Russia, and the Caucasus Region.

**IUCN status.** – Vulnerable under criterion B2ab(ii,iii,iv).

**Remarks.** – New species for Corsica.

*Callicera fagesii* Guérin-Méneville, 1844

**Material examined.** – 3 females.

**Distribution.** – Central and Southern Europe, especially in the Mediterranean Basin.

**IUCN status.** – Endangered under criterion B2ab(ii,iii,iv).

**Remarks.** – New species for Corsica.

*Callicera macquartii* Rondani, 1844

**Material examined.** – 2 females.

**Distribution.** – Central and Southern Europe, especially in the Mediterranean Basin

**IUCN status.** – Endangered under criterion B2ab(ii,iii,iv).

**Remarks.** – New species for Corsica.

*Ceriana conopoides* (Linnaeus, 1758)

**References.** – BECKER *et al.* (1910) as *Ceria conopoides*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male, 2 females.

**Distribution.** – Europe, Northern Africa, Russia (including Central and Eastern parts), and China.

**IUCN status.** – Least Concern.

*Ceriana vespiformis* (Latreille, 1809)

**References.** – BECKER *et al.* (1910) as *Ceria vespiformis*; DIRICKX (1994); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 16 males, 9 females.

**Distribution.** – Mediterranean Basin, including Middle East, the Balkan Peninsula up to Romania.

**IUCN status.** – Least Concern.

*Chalcosyrphus (Xylotina) nemorum* (Fabricius, 1805)

**References.** – KUNTZE (1913) as *Xylota nemorum*; DIRICKX (1994).

**Material examined.** – 19 males, 14 females.

**Distribution.** – British Isles, Europe, Russia, and Japan; also from Alaska to Nova Scotia and south to California in the Nearctic.

**IUCN status.** – Least Concern.

**Remarks.** – This is the second published record of this species from Corsica; being the first 110 years old. In recent literature, *C. nemorum* is not listed from Corsica (SPEIGHT, 2018; SPEIGHT *et al.*, 2018, 2020).

*Chalcosyrphus (Xylotodes) piger* (Fabricius, 1794)

**References.** – BIGOT (1861) and VILLENEUVE (1912) as *Xylota fulviventris* Bigot, 1861; BECKER *et al.* (1910) as *Xylota fulviventris* and *X. nigerrima* Becker, 1910; KUNTZE (1913) as *Xylota nigerrima*; VAN DER GOOT (1961) as *Zelima fulviventris*; HIPPA (1968) as *Xylotomima piger* because he considered *Xylotomima fulviventris* as a junior synonym of *X. piger*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 2 males.

**Distribution.** – Europe, Russia (including Russian Far East); and from British Columbia to Quebec, south to California, Mexico and Florida in the Nearctic.

**IUCN status.** – Least Concern.

***Cheilosia (Cheilosia) aerea*** Dufour, 1848

**References.** – BECKER *et al.* (1910) as *Chilosia zetterstedti* Becker, 1894; DIRICKX (1994) as *Cheilosia zetterstedti*; SPEIGHT *et al.* (2020).

**Material examined.** – 1 male, 2 females.

**Distribution.** – Central and Southern Europe, European parts of Russia, and Caucasus Region.

**IUCN status.** – Least Concern.

***Cheilosia (Cheilosia) proxima*** (Zetterstedt, 1843)

**Material examined.** – 2 females.

**Distribution.** – British Isles, Europe, Russia (including Central and Eastern part of Russia).

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Cheilosia (Convocheila) laticornis*** Rondani, 1857

**References.** – BECKER *et al.* (1910) as *Chilosia latifacies* Loew, 1857; DIRICKX (1994) as *Cheilosia latifacies*; SPEIGHT *et al.* (2020).

**Material examined.** – 1 male, 1 female.

**Distribution.** – Europe, south European Russia, Ukraine, the Caucasus Region, Kirghizstan, Afghanistan, Turkey, Israel, and Northern Africa.

**IUCN status.** – Least Concern.

***Cheilosia (Eucartosyrphus) scutellata*** (Fallén, 1817)

**References.** – KUNTZE (1913) as *Chilosia scutellata*; DIRICKX (1994); LEBARD *et al.* (2019); SPEIGHT *et al.* (2020).

**Material examined.** – 11 males, 22 females.

**Distribution.** – Europe, Turkey, Northern Africa, Russia (including the Russian Far East).

**IUCN status.** – Least Concern.

***Chrysotoxum bicinctum*** (Linnaeus, 1758)

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 6 males, 5 females.

**Distribution.** – Europe, Northern Africa, Turkey, Caucasus Region, and European parts of Russia eastwards until Central Siberia.

**IUCN status.** – Least Concern.

***Chrysotoxum cisalpinum*** Rondani, 1845

**References.** – SCHINER (1857); BECKER *et al.* (1910); VAN DER GOOT (1961); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male, 3 females.

**Distribution.** – Mediterranean Europe, central France, Bulgaria, Tajikistan, and Uzbekistan.

**IUCN status.** – Vulnerable under criterion B2ab(iii).

***Chrysotoxum elegans* Loew, 1841**

**References.** – BECKER *et al.* (1910); SPEIGHT *et al.* (2020).

**Material examined.** – 5 males, 1 female.

**Distribution.** – Europe, European parts of Russia, Turkey, and the Caucasus Region.

**IUCN status.** – Near Threatened under criterion B2ab(ii,iii,iv).

***Chrysotoxum intermedium* Meigen, 1822**

**References.** – SCHINER (1857); KUNTZE (1913); DIRICKX (1994).

**Distribution.** – Unclear.

**IUCN status.** – Least Concern.

**Remarks.** – The species name *C. intermedium* very likely comprises more than one taxon (SPEIGHT, 2018), and the species limits between *C. intermedium* and *C. lessonae* Giglio-Tos, 1890 are not clear yet (SPEIGHT & LEBARD, 2022). We identified the studied specimens using SPEIGHT & SARTHOU (2017) and named them *C. intermedium* A and *C. intermedium* B (see below). Following SPEIGHT & LEBARD (2022), the species concept of *C. intermedium* A corresponds globally to *Chrysotoxum lessonae* in SOMMAGGIO (2001) and VUJIĆ *et al.* (2017) and to the *C. lessonae* aggregate of SPEIGHT & LEBARD (2022). Similarly, *C. intermedium* B corresponds globally to *Chrysotoxum intermedium* in SOMMAGGIO (2001) and VUJIĆ *et al.* (2017) and to the *C. intermedium* aggregate of SPEIGHT & LEBARD (2022). We opted to name these two taxa *intermedium* A and *intermedium* B because the use of *C. lessonae* or *C. lessonae* aggregate would imply a new species record for Corsica; and although we are sure that both taxa are present in Corsica, we do not think it is necessary to mention *C. lessonae* from Corsica until the taxonomy and the nomenclature are revised.

***Chrysotoxum intermedium* A**

**References.** – SPEIGHT *et al.* (2018).

**Material examined.** – 4 males, 3 females.

**Remarks.** – See Remarks under *C. intermedium*.

***Chrysotoxum intermedium* B**

**References.** – LEBARD *et al.* (2019).

**Material examined.** – 14 males, 15 females.

**Remarks.** – See Remarks under *C. intermedium*.

***Chrysotoxum octomaculatum* Curtis, 1837**

**Material examined.** – 2 males, 2 females.

**Distribution.** – England, Central and Southern Europe, southern parts of Russia, Armenia, and Kazakhstan.

**IUCN status.** – Near Threatened.

**Remarks.** – New species for Corsica.

***Claussenia hispanica* (Strobl, 1909)**

**References.** – SPEIGHT *et al.* (1998) as *Heringia hispanica*; BIRTELE (2011) as *Heringia (Neocnemodon) hispanica*; SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 7 females.

**Distribution.** – Mediterranean Europe.

**IUCN status.** – Endangered under criterion B2ab(iii).

***Dasysyrphus albostriatus* (Fallén, 1817)**

**References.** – KUNTZE (1913) as *Syrphus albostriatus*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 6 males, 2 females.

**Distribution.** – British Isles, Europe, Northern Africa, Turkey, Caucasus Region, Russia, and Japan.

**IUCN status.** – Least Concern.

***Dasysyrphus pinastri* (De Geer, 1776)**

**Material examined.** – 1 male, 2 females.

**Distribution.** – British Isles, Europe, Turkey, and Russia (eastwards to Yakutia).

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Didea fasciata* Macquart, 1834**

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 2 males, 2 females.

**Distribution.** – British Isles, Europe, Russia (including Russian Far East), China, and Japan; India and Taiwan in the Indomalayan Region.

**IUCN status.** – Least Concern.

***Doros destillatorius* Mik, 1885**

**References.** – CORNUEL-WILLERMOZ (2021).

**Material examined.** – 1 female.

**Distribution.** – Mediterranean Europe, Bulgaria, Romania, Turkey and Crimea.

**IUCN status.** – Endangered under criterion B2ab(iii).

**Remarks.** – Second report of this species from Corsica.

***Epistrophe eligans* (Harris, 1780)**

**References.** – BECKER *et al.* (1910) as *Syrphus bifasciatus* Fabricius, 1794; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 4 males, 6 females.

**Distribution.** – British Isles, Europe, European parts of Russia, Turkey, and Caucasus Region.

**IUCN status.** – Least Concern.

**Remarks.** – All studied specimens belong to *Epistrophe eligans* var. *trifasciata* (Strobl, 1898), except for a male that belongs to the nominate subspecies.

***Epistrophe nitidicollis* (Meigen, 1822)**

**References.** – BECKER *et al.* (1910) as *Syrphus nitidicollis*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 4 females.

**Distribution.** – British Isles, Europe, Russia, Caucasus Region, and Japan; from Alaska south to California and South Carolina in the Nearctic Region.

**IUCN status.** – Least Concern.

***Episyphus balteatus* (De Geer, 1776)**

**References.** – BECKER *et al.* (1910) as *Syrphus balteatus*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 29 males, 26 females, 2 unknown.

**Distribution.** – British Isles, Europe, Canary Islands, Azores, Northern Africa, Russia, Caucasus Region, and Japan.

**IUCN status.** – Least Concern.

**Remarks.** – The two specimens with unknown sex are field observations.

***Eristalinus (Eristalinus) megacephalus* (Rossi, 1794)**

**References.** – DIRICKX (1994); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 3 females.

**Distribution.** – Mediterranean Europe, Northern Africa, Turkey, and Afrotropical Region.

**IUCN status.** – Least Concern.

***Eristalinus (Eristalinus) sepulchralis* (Linnaeus, 1758)**

**References.** – KUNTZE (1913); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 14 males, 13 females.

**Distribution.** – British Isles, Europe, Northern Africa, Turkey, Russia, China, Japan, and India.

**IUCN status.** – Least Concern.

***Eristalinus (Eristalodes) taeniops* (Wiedemann, 1818)**

**References.** – BECKER *et al.* (1910) as *Eristalis taeniops*; KUNTZE (1913) as *Eristalodes taeniops*; DIRICKX (1994); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 1 male, 1 unknown.

**Distribution.** – Mediterranean Europe, Middle East, Turkey, Northern Africa, Caucasus Region, Canary Islands, Afrotropical Region and parts of the Indomalayan Region.

**IUCN status.** – Least Concern.

**Remarks.** – The specimen with unknown sex is a field observation.

***Eristalinus (Lathyrophthalmus) aeneus* (Scopoli, 1763)**

**References.** – BECKER *et al.* (1910) as *Eristalis aeneus*; VAN DER GOOT (1961) as *Lathyrophthalmus aeneus*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 10 males, 8 females.

**Distribution.** – Cosmopolitan.

**IUCN status.** – Least Concern.

***Eristalis (Eoseristalis) arbustorum*** (Linnaeus, 1758)

**References.** – KUNTZE (1913); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 12 males, 13 females.

**Distribution.** – Throughout the Palaearctic Region, including Northern Africa; from Wisconsin to Labrador and south to Kansas and South Carolina in the Nearctic Region; and in northern India.

**IUCN status.** – Least Concern.

***Eristalis (Eoseristalis) pertinax*** (Scopoli, 1763)

**References.** – KUNTZE (1913); DIRICKX (1994); SPEIGHT *et al.* (2020).

**Material examined.** – 5 males.

**Distribution.** – British Isles, Europe, European parts of Russia, and Turkey.

**IUCN status.** – Least Concern.

***Eristalis (Eoseristalis) similis*** Fallén, 1817

**References.** – KUNTZE (1913) and DIRICKX (1994) as *Eristalis pratorum* Meigen, 1822; SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 9 males, 12 females.

**Distribution.** – England, Europe, Mediterranean Basin, Turkey, and Russia.

**IUCN status.** – Least Concern.

***Eristalis (Eristalis) tenax*** (Linnaeus, 1758)

**References.** – SCHINER (1857); BECKER *et al.* (1910) as *Eristalis hortorum* Meigen, 1822, and *E. tenax*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 25 males, 7 females.

**Distribution.** – Cosmopolitan.

**IUCN status.** – Least Concern.

***Eumerus amoenus*** Loew, 1848

**References.** – VAN DER GOOT (1961); DOCZKAL (1996).

**Material examined.** – 8 males, 4 females.

**Distribution.** – Central and Southern Europe, Mediterranean Basin, Canary Islands, Azores, Caucasus Region, and central parts of Asia (Kazakhstan, Turkestan, Tajikistan, and Mongolia).

**IUCN status.** – Least Concern.

***Eumerus argyropus*** Loew, 1848

**Material examined.** – 1 male.

**Distribution.** – Mediterranean Europe, Switzerland, Bulgaria, Romania, Ukraine, Caucasus Region, and Turkey.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Eumerus barbarus*** (Coquebert, 1804)

**References.** – BECKER *et al.* (1910) as *Eumerus iris* Loew, 1848 and *E. barbarus*; DIRICKX (1994); SPEIGHT (2014, 2018, 2020b); SPEIGHT & LEBARD (2020); SPEIGHT *et al.* (2020).

**Material examined.** – 6 males, 2 females.

**Distribution.** – Mediterranean Basin.

**IUCN status.** – Least Concern.

***Eumerus basalis* Loew, 1848**

**References.** – BECKER *et al.* (1910); VAN DER GOOT (1961); DIRICKX (1994); SPEIGHT & SARTHOU (2006); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male.

**Distribution.** – Mediterranean Europe, Bulgaria, Romania, Ukraine, Turkey, and Iran.

**IUCN status.** – Least Concern.

***Eumerus consimilis* Šimić & Vujić, 1996**

**Material examined.** – 1 male.

**Distribution.** – Unknown, but confirmed from Portugal, Croatia, France, and Sardinia.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Eumerus emarginatus* Loew, 1848**

**References.** – CORNUEL-WILLERMOZ *et al.* (2023).

**Material examined.** – 1 male.

**Distribution.** – Italy and Greece, including Peloponnesus and Aegean islands.

**IUCN status.** – Not Evaluated.

**Remarks.** – CORNUEL-WILLERMOZ *et al.* (2023) reported the species for the first time from Corsica including the male collected in the surveys of the *Our Planet Reviewed in Corsica 2019-2021*.

***Eumerus flavitarsis* Zetterstedt, 1843**

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT (2014, 2018, 2020b).

**Material examined.** – 13 males, 11 females.

**Distribution.** – Europe, Russia, and Japan.

**IUCN status.** – Least Concern.

***Eumerus niehuisi* Doczkal, 1996**

**References.** – DOCZKAL (1996); SPEIGHT *et al.* (1998, 2018, 2020); SPEIGHT (2014, 2018, 2020b).

**Material examined.** – 6 males, 2 females.

**Distribution.** – Corsica.

**IUCN status.** – Endangered under criterion B1ab(iii)+2ab(iii).

**Remarks.** – Endemic species from Corsica. RICARTE *et al.* (2012) reported two *Eumerus* males identified as *E. niehuisi* from the island of Lesvos (Greece) and they referred to the male genitalia and the antennal shape to ground their determination. GRKOVIĆ *et al.* (2015) provided new records of *Eumerus* specimens identified as *E. niehuisi* from the Greek islands of Chios, Lesvos, and Samos, and they established their determination by comparing the Greek material with the type material of *E. niehuisi*. Later, CHRONI *et al.* (2018) stated that the

material identified as *E. niehuisi* in Grković *et al.* (2015) belong to *Eumerus crassus* Grković, Vujić & Radenković in Grković *et al.* (2015). In their *Atlas of the Hoverflies of Greece*, VUJIĆ *et al.* (2020a) did not list *E. niehuisi* for Greece, neither MILIČIĆ & GRKOVIĆ (2021) in their assessment for the European Red List of Hoverflies. Thus, we interpret that *E. niehuisi* is considered absent from Greece by VUJIĆ *et al.* (2020a).

CHRONI *et al.* (2018) also mentioned Corsica and Sardinia for the distributional range of *E. niehuisi* referring to DOCZKAL (1996) as the source for such statement, but DOCZKAL (1996) did not study (neither mentioned) any individual from Sardinia. Thus, *E. niehuisi* does not occur on Sardinia based on the current evidence.

### *Eumerus obliquus* (Fabricius, 1805)

**References.** – SPEIGHT *et al.* (1998); BIRTELE (2011); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 3 males.

**Distribution.** – Western Mediterranean Basin (including islands), Canary Islands, Afro-tropical Region (including the Mascarene Islands and Madagascar), Yemen, Socotra, and Australia.

**IUCN status.** – Least Concern.

### *Eumerus ornatus* Meigen, 1822

**Material examined.** – 2 males.

**Distribution.** – England, Central and Southern Europe, Romania, Ukraine, and Caucasus Region.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica. We believe that the specimens reported by BECKER *et al.* (1910) and DIRICKX (1994) as *Eumerus lucidus* Loew, 1848 from Corsica may refer to this taxon, but we did not study this old material.

### *Eumerus pulchellus* Loew, 1848

**References.** – BECKER *et al.* (1910); VAN DER GOOT (1961); DIRICKX (1994); DOCZKAL (1996); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 64 males, 9 females, 1 intersex.

**Distribution.** – Southern Europe and Mediterranean Basin.

**IUCN status.** – Least Concern.

### *Eumerus* sp.

**Material examined.** – 14 females.

**Remarks.** – All these female specimens belong to several different species, but we cannot properly identify them at this moment.

### *Eumerus sulcitibius* Rondani, 1868

**References.** – DIRICKX (1994); SPEIGHT *et al.* (1998, 2018, 2020); BIRTELE (2011); SPEIGHT (2014, 2018, 2020b).

**Material examined.** – 5 males, 2 females.

**Distribution.** – Southern Europe, Turkey, eastwards to Azerbaijan.

**IUCN status.** – Least Concern.

***Eumerus vandenberghhei*** Doczkal, 1996

**References.** – DOCZKAL (1996); SPEIGHT *et al.* (1998, 2018, 2020); BIRTELE (2011); SPEIGHT (2014, 2018, 2020b); GRKOVIĆ *et al.* (2017).

**Material examined.** – 1 male.

**Distribution.** – Corsica and Sardinia.

**IUCN status.** – Endangered under criterion B1ab(iii)+2ab(iii).

***Eupeodes (Eupeodes) corollae*** (Fabricius, 1794)

**References.** – SCHINER (1857) as *Syrphus corollae*; KUNTZE (1913) as *Syrphus corollae*; DIRICKX (1994) as *Metasyrphus corollae*; SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 16 males, 46 females.

**Distribution.** – Palaearctic and Afrotropical Regions, also Taiwan.

**IUCN status.** – Least Concern.

***Eupeodes (Eupeodes) latifasciatus*** (Macquart, 1829)

**Material examined.** – 6 males, 2 females.

**Distribution.** – British Isles, Europe, Northern Africa, Turkey, and Russia (including Russian Far East); India; and from Alaska south to California and Texas in the Nearctic Region.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Eupeodes (Eupeodes) lucasi*** (Marcos-García & Láska, 1983)

**References.** – MARCOS-GARCÍA *et al.* (2000); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 17 males, 5 females.

**Distribution.** – Central and Southern Europe.

**IUCN status.** – Least Concern.

***Eupeodes (Eupeodes) luniger*** (Meigen, 1822)

**References.** – BECKER *et al.* (1910) as *Syrphus luniger*; DIRICKX (1994) as *Metasyrphus luniger*.

**Material examined.** – 5 males, 4 females.

**Distribution.** – British Isles, Europe, Madeira, Northern Africa, Turkey, Russia (including Russian Far East), and Japan; also in Northern India.

**IUCN status.** – Least Concern.

**Remarks.** – This is the second published record of this species from Corsica; being the first 110 years old. In recent literature, *E. luniger* is not listed from Corsica (SPEIGHT, 2018; SPEIGHT *et al.*, 2018, 2020).

***Eupeodes (Eupeodes) nuba*** (Wiedemann, 1830)

**References.** – SPEIGHT *et al.* (2020).

**Material examined.** – 1 male, 4 females.

**Distribution.** – Canary Isles, Mediterranean Basin, Switzerland, Romania, Caucasus Region, south-western parts of Asia (Uzbekistan, Kirghizstan, Tajikistan), Afghanistan and Mongolia; and from Ethiopia south to South Africa in the Afrotropical Region.

**IUCN status.** – Least Concern.

***Eupeodes (Eupeodes) vandergooti*** (Dušek & Láska, 1973)

**References.** – DUŠEK & LÁSKA (1973) as *Metasyrphus vandergooti*; DIRICKX (1994) as *Metasyrphus vandergooti*; SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male, 7 females.

**Distribution.** – Corsica.

**IUCN status.** – Endangered under criterion B1ab(iii,iv)+2ab(iii,iv).

**Remarks.** – Endemic from Corsica. SPEIGHT (2020b) listed this species from Italy, but MAZÁNEK *et al.* (2021) considered uncertain the presence of *E. vandergooti* from mainland France, mainland Italy and Sardinia.

***Fagisyrphus cinctus*** (Fallén, 1817)

**References.** – BECKER *et al.* (1910) as *Syrphus cinctus*; DIRICKX (1994) as *Melangyna cincta*; SPEIGHT *et al.* (2020) as *Meligramma cincta*.

**Material examined.** – 2 females.

**Distribution.** – Europe, Crimea, European parts of Russia, Caucasus Region, and Turkey.

**IUCN status.** – Least Concern.

***Ferdinandea cuprea*** (Scopoli, 1763)

**Material examined.** – 1 male.

**Distribution.** – British Isles, Europe, Northern Africa, Turkey, Russia, and Japan.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Helophilus (Helophilus) pendulus*** (Linnaeus, 1758)

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 11 males, 5 females.

**Distribution.** – British Isles, Europe, and Russia (including the Pacific coast).

**IUCN status.** – Least Concern.

***Helophilus (Helophilus) trivittatus*** (Fabricius, 1805)

**References.** – KUNTZE (1913) as *Heliophilus bivittatus* Fabricius [misspelling]; SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male, 1 female.

**Distribution.** – British Isles, Europe, eastwards through Russian Federation to the Pacific, including Iran and Afghanistan.

**IUCN status.** – Least Concern.

***Heringia heringi*** (Zetterstedt, 1843)

**References.** – BECKER *et al.* (1910) as *Pipizella heringi*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 4 males, 1 female.

**Distribution.** – British Isles, Europe, Turkey, European parts of Russia, and Mongolia.

**IUCN status.** – Least Concern.

***Lapposyrphus lapponicus* (Zetterstedt, 1838)**

**Material examined.** – 2 males, 1 female.

**Distribution.** – British Isles, Europe, Turkey, Russia (to the Pacific coast), Japan, Iceland, and Greenland. Its presence in North America needs re-assessment.

**IUCN status.** – Least Concern.

**Remarks.** – New species and new genus for Corsica.

***Lejogaster tarsata* (Meigen, 1822)**

**References.** – BECKER *et al.* (1910) as *Liogaster splendida* (Meigen, 1822); DIRICKX (1994); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 4 males, 15 females.

**Distribution.** – British Isles, Europe, Russia (eastwards until the Pacific coast), Middle East, and Asia (Afghanistan, Uzbekistan, Tajikistan, Kirghizia, Turkmenia, Kazakhstan, Mongolia).

**IUCN status.** – Least Concern.

***Lejops vittatus* (Meigen, 1822) (fig. 2-4)**

**Material examined.** – 3 males, 9 females.



**Fig. 2-4.** – *Lejops vittatus* (Meigen, 1822), new species and new genus records from Corsica. – 2, Sampling locality of *L. vittatus*, marais de Péri. – 3, Male resting. – 4, Female. © T. Lebard.

**Distribution.** – England, Europe, Russian Federation (eastwards to the Pacific coast, including middle Asia).

**IUCN status.** – Vulnerable under criterion A3c.

**Remarks.** – New species and new genus for Corsica.

***Mallota cimbiciformis* (Fallén, 1817)**

**References.** – BECKER *et al.* (1910) as *Mallota cymbiciformis* Fallén [misspelling]; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male, 1 female.

**Distribution.** – British Isles, Europe, Northern Africa, Russia (to central Siberia), and Iran.

**IUCN status.** – Least Concern.

***Melangyna compositarum* (Verrall, 1873)**

**Material examined.** – 1 female.

**Distribution.** – British Isles, Europe, Russia (to the Russian Far East), and Japan; and from Alaska south through the Rocky Mountains to New Mexico in the Nearctic Region.

**IUCN status.** – Least Concern.

**Remarks.** – New species and new genus for Corsica.

***Melanostoma mellinum* (Linnaeus, 1758)**

**References.** – BECKER *et al.* (1910); VAN DER GOOT (1961); DIRICKX (1994); LEBARD *et al.* (2019); SPEIGHT *et al.* (2020).

**Material examined.** – 42 males, 53 females.

**Distribution.** – Northern Africa, Palaearctic Region, and from Alaska to Quebec and south to Washington in the Nearctic Region.

**IUCN status.** – Least Concern.

***Melanostoma scalare* (Fabricius, 1794)**

**References.** – BECKER *et al.* (1910); DIRICKX (1994); LEBARD *et al.* (2019); SPEIGHT *et al.* (2020).

**Material examined.** – 94 males, 70 females.

**Distribution.** – British Isles, Europe, Northern Africa, Russia, Japan, and China. records from the Afrotropical, Australasian and Indomalayan regions need re-assessment.

**IUCN status.** – Least Concern.

***Meliscaeva auricollis* (Meigen, 1822)**

**References.** – BECKER *et al.* (1910) as *Syrphus auricollis* and *Syrphus maculicornis* var. *nigritibius* Rondani, 1857; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 22 males, 12 females, 1 unknown.

**Distribution.** – British Isles, Europe, Canary Islands, Northern Africa, Middle East, Caucasus Region, and European parts of Russia.

**IUCN status.** – Least Concern.

**Remarks.** – The specimen with unknown sex is a field observation.

***Meliscaeva cinctella* (Zetterstedt, 1843)**

**References.** – BECKER *et al.* (1910) as *Syrphus cinctellus*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 16 males, 7 females.

**Distribution.** – British Isles, Europe, Northern Africa, Turkey, Caucasus Region, Russia (including the Russian Far East), Japan, and China; Alaska south to California and Colorado in the Nearctic Region.

**IUCN status.** – Least Concern.

***Merodon aff. aureus***

**References.** – SPEIGHT & LANGLOIS (2020) as *Merodon* species C; JANKOVIĆ & RADENKOVIĆ (2021) as *Merodon aerarius* Rondani, 1857.

**Material examined.** – 22 males, 17 females.

**Distribution.** – Austria, France (including Corsica), Italy (including Sardinia), Balkan Peninsula, and Romania.

**IUCN status.** – Least Concern.

**Remarks.** – VUJIĆ *et al.* (2021: Supplementary file S4) listed *Merodon aerarius* as a valid species of the *cinereus* subgroup within the *aureus* group. This taxon was named ‘species C’ in SPEIGHT & LANGLOIS (2020), and JANKOVIĆ & RADENKOVIĆ (2021) referred to it as *Merodon aerarius*. Our specimens belong to the same taxon as the one reported by SPEIGHT & LANGLOIS (2020) and JANKOVIĆ & RADENKOVIĆ (2021) from Corsica, whose name should be *Merodon aerarius* (Ante VUJIĆ, pers. comm.). We hesitate to name it *M. aerarius* in the present work until a ground-based taxonomic decision is taken (VUJIĆ *et al.*, in prep.).

***Merodon avidus* (Rossi, 1790)**

**References.** – SCHINER (1857) as *Merodon spinipes* (Fabricius, 1794) and *M. avidus*; BECKER *et al.* (1910) as *Merodon spinipes*; KUNTZE (1913); VAN DER GOOT (1961) as *Lampetia spinipes*; HURKMAN (1993); DIRICKX (1994); POPOVIĆ *et al.* (2015); SPEIGHT & LEBARD (2020); SPEIGHT *et al.* (2020).

**Material examined.** – 46 males, 10 females.

**Distribution.** – Mediterranean Europe and Romania.

**IUCN status.** – Least Concern.

***Merodon clavipes* (Fabricius, 1781)**

**References.** – SCHINER (1857); BECKER *et al.* (1910); DIRICKX (1994); MARCOS-GARCÍA *et al.* (2007); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 2 females.

**Distribution.** – Central and Southern Europe, Romania, Ukraine, Turkey, and Northern Africa.

**IUCN status.** – Least Concern.

***Merodon equestris* (Fabricius, 1794)**

**References.** – BECKER *et al.* (1910) as *Merodon equestris* Fbr. var. *nigrithorax* Bezzi, 1900; DIRICKX (1994).

**Material examined.** – 4 males.

**Distribution.** – British Isles, Europe, Northern Africa, Russia (including Russian Far East), and Japan; also in New Zealand (introduced), and from British Columbia south to California in the Nearctic Region.

**IUCN status.** – Least Concern.

**Remarks.** – This is the second published record of this species from Corsica; being the first 110 years old. In recent literature, *M. equestris* is not listed from Corsica (SPEIGHT, 2018; SPEIGHT *et al.*, 2018, 2020).

***Merodon femoratus* Sack, 1913**

**References.** – HURKMANS (1993); DIRICKX (1994); SPEIGHT (2018); SPEIGHT *et al.* (2018, 2020); SPEIGHT & LANGLOIS (2020).

**Material examined.** – 2 males.

**Distribution.** – Mediterranean Basin.

**IUCN status.** – Least Concern.

***Merodon funestus* (Fabricius, 1794)**

**Material examined.** – 1 male, 2 females.

**Distribution.** – Mediterranean Basin (except Northern Africa).

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Merodon geniculatus* Strobl, 1909**

**References.** – DIRICKX (1994); SPEIGHT *et al.* (1998, 2018, 2020); MARCOS-GARCÍA *et al.* (2007); SPEIGHT (2014, 2018, 2020b); RICARTE *et al.* (2017); SPEIGHT & LANGLOIS (2020).

**Material examined.** – 4 males.

**Distribution.** – Mediterranean Basin.

**IUCN status.** – Near Threatened under criterion B2ab(iii).

***Merodon minutus* Strobl, 1893**

**References.** – DIRICKX (1994); SPEIGHT (2018); SPEIGHT *et al.* (2018, 2020); SPEIGHT & LANGLOIS (2020); AČANSKI *et al.* (2022).

**Material examined.** – 10 males, 12 females.

**Distribution.** – It needs a re-assessment due to recent nomenclatural changes (AČANSKI *et al.*, 2022), but it is known from Corsica, Balkan Peninsula, Sardinia, Sicily, and Greece (East Aegean Is., mainland, and Crete).

**IUCN status.** – Least Concern.

***Merodon natans* (Fabricius, 1794)**

**Material examined.** – 1 female.

**Distribution.** – Mediterranean Europe, Bulgaria, Israel, and Caucasus Region.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Merodon rubidiventris* Costa, 1884**

**References.** – HURKMANS (1993), SPEIGHT *et al.* (1998, 2018, 2020), SPEIGHT (2014, 2018, 2020b) and LEBARD *et al.* (2019) as *Merodon mariae* Hurkmans, 1993; SPEIGHT & LANGLOIS (2020); VUJIĆ *et al.* (2020b).

**Material examined.** – 2 males, 3 females.

**Distribution.** – Corsica and Sardinia.

**IUCN status.** – Vulnerable under criterion B2ab(iii).

***Merodon trochantericus*** Costa, 1884

**References.** – VILLENEUVE (1909) as *Merodon podagricus* Villeneuve, 1909; BECKER *et al.* (1910) as *Merodon podagricus*; DIRICKX (1994); MARCOS-GARCÍA *et al.* (2007); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 3 males.

**Distribution.** – France, Spain, Italy, Corsica, Sardinia, and Corfu (Greece).

**IUCN status.** – Least Concern.

***Mesembrius peregrinus*** (Loew, 1846)

**Material examined.** – 6 males, 3 females.

**Distribution.** – Central and Southern Europe, Israel, Ukraine, Caucasus Region, Russia, Japan, and China.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Milesia crabroniformis*** (Fabricius, 1775)

**References.** – KUNTZE (1913); DIRICKX (1994); SARTHOU *et al.* (2004); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 2 males, 2 females.

**Distribution.** – Mediterranean Europe, Portugal, Madeira, Switzerland, Bulgaria, Hungary, and Turkey.

**IUCN status.** – Least Concern.

***Milesia semiluctifera*** (Villers, 1789)

**References.** – VAN DER GOOT (1961); DIRICKX (1994); SPEIGHT & LEBARD (2020); SPEIGHT *et al.* (2020).

**Material examined.** – 3 males.

**Distribution.** – Mediterranean Basin, Switzerland, Romania, Ukraine, Caucasus Region, eastwards to Turkmenistan.

**IUCN status.** – Least Concern.

***Myathropa florea*** (Linnaeus, 1758)

**References.** – SCHINER (1857) as *Helophilus floreus*; BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 12 males, 7 females.

**Distribution.** – Palaearctic Region.

**IUCN status.** – Least Concern.

***Myolepta dubia*** (Fabricius, 1805) (fig. 7-8)

**References.** – KUNTZE (1913) as *Myiolepta luteola* (Gmelin, 1790); DIRICKX (1994) as *Myolepta luteola*; SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 8 males, 1 female.

**Distribution.** – British Isles, Europe, and European parts of Russia.

**IUCN status.** – Least Concern.

***Myolepta nigritarsis* Coe, 1957 (fig. 5-6)**

**Material examined.** – 1 male.

**Distribution.** – Southern France, Italy, Austria, Hungary, Romania, Greece, Crete, Balkan Peninsula, southern parts of European Russia, Turkey, Azerbaijan, and Armenia.


**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Myolepta* sp.**

**Material examined.** – 2 females.

**Remarks.** – We are not sure about the species identification of these two females. They were collected together with *Myolepta dubia* males, but we key them out to *Myolepta potens*



**Fig. 5-8.** – *Myolepta* spp. – 5-6, Sampling locality of *Myolepta nigritarsis* Coe and *Mallota cimbiciformis* (Fallén), 1 km west of Valavo. – 7-8, Male of *Myolepta dubia* (Fabricius). © T. Lebard.

in the identification key by REEMER *et al.* (2004). We need a further comparison with more material of *M. dubia* and, maybe, a molecular study.

***Neoascia (Neoascia) podagraria* (Fabricius, 1775)**

**References.** – BECKER *et al.* (1910) as *Ascia podagraria*; KUNTZE (1913) as *Ascia floralis* Meigen, 1822; DIRICKX (1994) as *Neoascia floralis* and *N. podagraria*; SPEIGHT *et al.* (2020).

**Material examined.** – 2 females.

**Distribution.** – British Isles, Europe, Mediterranean Basin, Madeira, and Russia (eastwards to Cis-Baikal).

**IUCN status.** – Least Concern.

***Neoascia (Neoascia) tenur* (Harris, 1780)**

**Material examined.** – 3 males, 1 female.

**Distribution.** – Western Palaearctic (including Iceland), eastwards until most of Siberia.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Neoascia (Neoasciella) interrupta* (Meigen, 1822)**

**Material examined.** – 19 males, 8 females.

**Distribution.** – Northern and Central Europe, England, European parts of Russia (until Siberia), and the Caucasus Region.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Neocnemodon brevidens* (Egger, 1865)**

**Material examined.** – 4 males, 4 females.

**Distribution.** – England, Central Europe (northwards until Latvia), Romania, Ukraine, Russia (across Siberia to the Pacific coast), and China.

**IUCN status.** – Least Concern.

**Remarks.** – New species and new genus for Corsica.

***Neocnemodon latitarsis* (Egger, 1865)**

**Material examined.** – 22 males, 16 females.

**Distribution.** – Britain, Europe, and the Caucasus Region.

**IUCN status.** – Least Concern.

**Remarks.** – New species and new genus for Corsica.

***Neocnemodon* sp.**

**Material examined.** – 1 female.

**Remarks.** – The specimen is damaged and not well-preserved. It keys out to *Neocnemodon larusi* Vujić, 1999 using the identification key of BARTSCH (2009).

***Neocnemodon vitripennis* (Meigen, 1822)**

**Material examined.** – 3 females.

**Distribution.** – British Isles, Northern and Central Europe, Russia (until the Pacific coast), China, and Japan.

**IUCN status.** – Least Concern.

**Remarks.** – New species and new genus for Corsica.

***Paragus (Pandasyophthalmus) ascoensis*** Goedlin de Tiefenau & Lucas, 1981

**References.** – GOEDLIN DE TIEFENAU & LUCAS (1981); DIRICKX (1994); BIRTELE (2011); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 29 males.

**Distribution.** – Corsica and Sardinia.

**IUCN status.** – Vulnerable under criterion B2ab(iii).

***Paragus (Pandasyophthalmus) haemorrhous*** Meigen, 1822

**References.** – GOEDLIN DE TIEFENAU & LUCAS (1981); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 58 males, 2 females.

**Distribution.** – British Isles, Europe, Mediterranean Basin, Russia, Caucasus Region, Japan, and China; the Afrotropical Region; and from Yukon south to Costa Rica in the Americas.

**IUCN status.** – Least Concern.

***Paragus (Paragus) albifrons*** (Fallén, 1817)

**References.** – BECKER *et al.* (1910); GOEDLIN DE TIEFENAU & LUCAS (1981); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 2 males.

**Distribution.** – British Isles, Europe, Mediterranean Basin, Caucasus Region, Russia (until the Pacific coast), Mongolia, and China.

**IUCN status.** – Endangered under criterion B2ab(iii,iv).

***Paragus (Paragus) bicolor*** (Fabricius, 1794)

**References.** – KUNTZE (1913); GOEDLIN DE TIEFENAU & LUCAS (1981); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 24 males, 7 females.

**Distribution.** – Europe, Mediterranean Basin, Iran, Afghanistan, Russia, Mongolia, and China.

**IUCN status.** – Least Concern.

***Paragus (Paragus) bradescui*** Stănescu, 1981

**References.** – GOEDLIN DE TIEFENAU & LUCAS (1981) and DIRICKX (1994) as *Paragus antoinettae* Goedlin de Tiefenau & Lucas, 1981; SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male.

**Distribution.** – France, Portugal, Spain, Italy, Corsica, Sardinia, Sicily, Balkan Peninsula, Romania, Greece, Ukraine, Russia, Kyrgyzstan, Turkmenistan, and Tajikistan.

**IUCN status.** – Endangered under criterion B2ab(iii).

***Paragus (Paragus) pecchiolii*** Rondani, 1857

**References.** – SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 22 males, 3 females.

**Distribution.** – Europe, Mediterranean Basin, Turkey, Caucasus Region, and European parts of Russia.

**IUCN status.** – Least Concern.

***Paragus (Paragus) quadrifasciatus* Meigen, 1822**

**References.** – SCHINER (1857); GOELDLIN DE TIEFENAU & LUCAS (1981); DIRICKX (1994); SPEIGHT et al. (2018, 2020).

**Material examined.** – 5 males.

**Distribution.** – Southern Europe, France, Romania, Northern Africa, Turkey, Caucasus Region, Iran, Russia, middle Asia, and China.

**IUCN status.** – Least Concern.

***Paragus (Paragus) sexarciatus* Bigot, 1862**

**References.** – BIGOT (1862); BECKER et al. (1910); GOELDLIN DE TIEFENAU & LUCAS (1981); DIRICKX (1994); BIRTELE (2011); SPEIGHT (2014, 2018, 2020b); SPEIGHT et al. (2018, 2020); LEBARD et al. (2019).

**Material examined.** – 1 male.

**Distribution.** – Corsica, Sardinia, and Sicily.

**IUCN status.** – Vulnerable under criteria B1ab(iii)+2ab(iii).

***Paragus (Paragus) strigatus* Meigen, 1822**

**References.** – KUNTZE (1913); GOELDLIN DE TIEFENAU & LUCAS (1981); DIRICKX (1994); SPEIGHT et al. (2018, 2020).

**Material examined.** – 37 males, 7 females.

**Distribution.** – Portugal, Mediterranean Basin, Bulgaria, Romania, Ukraine, Western Russia to Kirghizia, Tajikistan, and Mongolia.

**IUCN status.** – Least Concern.

***Paragus* sp.**

**Material examined.** – 16 females.

**Remarks.** – The unnamed females may belong to several *Paragus* species, all of them to the subgenus *Pandasyophthalmus*, except for one female that belongs to the *bicolor* group of the subgenus *Paragus*.

***Parhelophilus frutetorum* (Fabricius, 1775)**

**Material examined.** – 2 males.

**Distribution.** – Europe, Caucasus Region, Russia, and China.

**IUCN status.** – Least Concern.

**Remarks.** – New species and new genus for Corsica.

***Parhelophilus versicolor* (Fabricius, 1794)**

**Material examined.** – 2 males, 6 females.

**Distribution.** – British Isles, Europe, Mediterranean Basin, Russia, and China.

**IUCN status.** – Least Concern.

**Remarks.** – New species (and new genus) for Corsica.

**Pelecocera (*Chamaesyphus*) *lusitanica*** (Mik, 1898)

**Material examined.** – 1 male, 1 female.

**Distribution.** – Europe and European parts of Russia.

**IUCN status.** – Near Threatened under criterion A3c.

**Remarks.** – New species for Corsica. MENGUAL *et al.* (2015b) synonymized *P. lusitanica* under *Pelecocera (Chamaesyphus) lugubris* Perris, 1839 without studying the type material, but VAN ECK & MENGUAL (2021) suggested to use *P. lusitanica* until a comprehensive molecular study can be carried out using specimens from Northern, Central and Southern Europe, including the Iberian Peninsula. LAIR *et al.* (2022) demonstrated that the name for the specimens identified as *P. lusitanica* from mainland France (all records are from the Atlantic coast) is *P. lugubris*. Thus, the question about the synonymy proposed by MENGUAL *et al.* (2015b) remains open and we decided to use *P. lusitanica* for the studied Corsican specimens as they do not originate from mainland France.

The male specimen used in the molecular study by VAN ECK & MENGUAL (2021) is the same male specimen reported here (unique identifier: LPRC2021-2417; COI sequence GenBank accession number: OK330482).

**Pelecocera (*Chamaesyphus*) *pruinosomaculata*** Strobl, 1906

**References.** – VAN ECK & MENGUAL (2021); LAIR *et al.* (2022).

**Material examined.** – 1 female.

**Distribution.** – Uncertain due to confusion with related species, but known from Portugal, Spain, southern France, Corsica, southern Italy, Cyprus, and Greece.

**IUCN status.** – Near Threatened under criterion A3c.

**Platycheirus (*Platycheirus*) *fulviventris*** (Macquart, 1829)

**References.** – BECKER *et al.* (1910) as *Platychirus fulviventris*; DIRICKX (1994); SPEIGHT *et al.* (2020).

**Material examined.** – 22 males, 10 females.

**Distribution.** – Europe, Morocco, Turkey, and Russia (to the Pacific coast).

**IUCN status.** – Least Concern.

**Platycheirus (*Platycheirus*) *muelleri*** Marcuzzi, 1941

**References.** – SPEIGHT *et al.* (1998, 2018, 2020); SPEIGHT (2014, 2018, 2020b).

**Material examined.** – 71 males, 64 females, 2 intersex.

**Distribution.** – Southern France, Corsica, Sardinia, northern Italy, and mainland Greece.

**IUCN status.** – Endangered under criterion B2ab(ii).

**Platycheirus (*Platycheirus*) *scutatus*** (Meigen, 1822)

**Material examined.** – 13 males, 20 females.

**Distribution.** – Europe, Turkey, Russia, Afghanistan, and China.

**IUCN status.** – Least Concern.

**Remarks.** – New species for Corsica.

***Platycheirus (Platycheirus) sp.***

**Material examined.** – 3 females.

**Remarks.** – The preservation condition of these three females is not optimal and their identification was not possible.

***Pyrophaena rosarum* (Fabricius, 1787)**

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 4 males, 4 females.

**Distribution.** – Holarctic, but not in northern Africa.

**IUCN status.** – Least Concern.

***Riponnensisia daccordii* (Claussen, 1991)**

**References.** – CLAUSSEN (1991) and DIRICKX (1994) as *Orthonevra daccordii*; SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 3 males.

**Distribution.** – Corsica.

**IUCN status.** – Critically Endangered under criteria B1ab(i,ii,iii)+2ab(i,ii,iii).

**Remarks.** – Endemic for Corsica. VAN STEENIS *et al.* (2021) indicated that the five females identified as *Orthonevra* sp. in BECKER *et al.* (1910) might refer to this taxon, but this is not confirmed. The same authors assessed this taxon as Critically Endangered and Possibly Extinct (PE), but our records from July 2022 indicate that there is still a population, which comes from a location (near Quenza) different from the type locality that VAN STEENIS *et al.* (2021) used in their assessment (between Tolla and Bastelica); the only known location for this species before our survey.

***Riponnensisia splendens* (Meigen, 1822)**

**References.** – BECKER *et al.* (1910) as *Chrysogaster splendens*; DIRICKX (1994) as *Orthonevra splendens*; SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 11 males, 7 females.

**Distribution.** – British Isles, Central and Southern Europe, Mediterranean Basin, Crimea, and the Caucasus Region.

**IUCN status.** – Least Concern.

***Scaeava (Scaeava) albomaculata* (Macquart, 1842)**

**References.** – SCHINER (1857) as *Syphus gemellarii* (Rondani, 1845); BECKER *et al.* (1910) as *Catabomba albomaculata*; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 2 males.

**Distribution.** – Mediterranean Basin, Canary Islands, Caucasus Region, southern Russia eastward to China and Mongolia.

**IUCN status.** – Least Concern.

***Scaeva (Semiscaeva) dignota* (Rondani, 1857)**

**References.** – DIRICKX (1994); SPEIGHT *et al.* (2020).

**Material examined.** – 26 males, 3 females.

**Distribution.** – England, Central and Southern Europe, Mediterranean Basin, and Canary Islands.

**IUCN status.** – Least Concern.

***Scaeva (Semiscaeva) mecogramma* (Bigot, 1860)**

**References.** – DIRICKX (1994); BIRTELE (2011); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male.

**Distribution.** – France, Portugal, Spain, Corsica, mainland Italy, Sardinia, Sicily, and Greece.

**IUCN status.** – Least Concern.

***Sericomyia silentis* (Harris, 1778)**

**References.** – KUNTZE (1913) as *Sericomyia borealis* (Fallén, 1816); DIRICKX (1994).

**Material examined.** – 1 female.

**Distribution.** – British Islands, Europe, Caucasus Region, and European parts of Russia.

**IUCN status.** – Least Concern.

**Remarks.** – This is the second published record of this species from Corsica; being the first 110 years old. In recent literature, *S. silentis* is not listed from Corsica (SPEIGHT, 2018; SPEIGHT *et al.*, 2018; SPEIGHT *et al.*, 2020).

***Sphaerophoria (Sphaerophoria) rueppellii* (Wiedemann, 1830)**

**References.** – BECKER *et al.* (1910) as *Sphaerophoria flavicauda* var. *nitidicollis* Zetterstedt, 1849; DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 17 males, 1 female.

**Distribution.** – Palaearctic and eastern parts of the Afrotropical Region.

**IUCN status.** – Least Concern.

***Sphaerophoria (Sphaerophoria) scripta* (Linnaeus, 1758)**

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 101 males, 50 females.

**Distribution.** – Palaearctic Region.

**IUCN status.** – Least Concern.

***Sphegina (Sphegina) clunipes* (Fallén, 1816)**

**References.** – BECKER *et al.* (1910); DIRICKX (1994); LEBARD *et al.* (2019); SPEIGHT *et al.* (2020).

**Material examined.** – 15 males, 10 females.

**Distribution.** – Western Palaearctic.

**IUCN status.** – Least Concern.

*Sphegina (Sphegina) elegans* Schummel, 1841

**References.** – DIRICKX (1994).

**Material examined.** – 8 males, 11 females.

**Distribution.** – British Isles, Europe, Caucasus Region, and European parts of Russia.

**IUCN status.** – Least Concern.

*Sphiximorpha subsessilis* (Illiger, 1807) (fig. 9-10)

**References.** – SCHINER (1857) as *Ceria subsessilis*; DIRICKX (1994); SPEIGHT *et al.* (2020).

**Material examined.** – 4 males.

**Distribution.** – Europe, Caucasus Region, and European parts of Russia

**IUCN status.** – Least Concern.

*Syritta flaviventris* Macquart, 1842

**References.** – VAN DER GOOT (1961) as *Syritta fasciata* (Wiedemann, 1830); VAN DER GOOT (1964) as *Syritta spinigera* Loew, 1848; DIRICKX (1994); SPEIGHT (2014, 2018, 2020b); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 1 male, 1 female.

**Distribution.** – Mediterranean Basin, Middle East, the Afrotropical Region, Central and South America, and Eastern Island.

**IUCN status.** – Least Concern.

*Syritta pipiens* (Linnaeus, 1758)

**References.** – SCHINER (1857); BECKER *et al.* (1910); KUNTZE (1913); VAN DER GOOT (1961); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 31 males, 19 females.



9



10

Fig. 9-10. – *Sphiximorpha subsessilis* (Illiger, 1807), collected 2 km south of Fromontica. © T. Lebard.

**Distribution.** – Holarctic Region.

**IUCN status.** – Least Concern.

***Syrphus ribesii* (Linnaeus, 1758)**

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 55 males, 14 females.

**Distribution.** – Holarctic Region.

**IUCN status.** – Least Concern.

***Syrphus torvus* Osten Sacken, 1875**

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2020).

**Material examined.** – 1 female.

**Distribution.** – Holarctic Region; and Taiwan, northern India, Nepal, and Thailand in the Indomalayan Region.

**IUCN status.** – Least Concern.

***Syrphus vitripennis* Meigen, 1822**

**References.** – KUNTZE (1913); DIRICKX (1994); SPEIGHT *et al.* (2020).

**Material examined.** – 14 males, 13 females.

**Distribution.** – Holarctic Region.

**IUCN status.** – Least Concern.

***Volucella zonaria* (Poda, 1761)**

**References.** – BECKER *et al.* (1910); VAN DER GOOT (1961) as *Volucella zonaria beckeri* Van der Goot, 1961; DIRICKX (1994); BIRTELE (2011); SPEIGHT (2014, 2018); LEBARD *et al.* (2019).

**Material examined.** – 12 males, 14 females.

**Distribution.** – British Isles, Central and Southern Europe, Russia, Iran, Mongolia, and China.

**IUCN status.** – Least Concern.

***Xanthandrus comtus* (Harris, 1780)**

**References.** – KUNTZE (1913); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 7 males, 3 females.

**Distribution.** – Palaearctic Region.

**IUCN status.** – Least Concern.

***Xanthogramma stackelbergi* Violovitsh, 1975**

**References.** – SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 4 males, 15 females.

**Distribution.** – Uncertain due to the confusion with other *Xanthogramma* species, but known from most Europe, England, Caucasus Region, and European parts of Russia.

**IUCN status.** – Least Concern.

**Xylota segnis** (Linnaeus, 1758)

**References.** – BECKER *et al.* (1910); SPEIGHT *et al.* (2018, 2020); LEBARD *et al.* (2019).

**Material examined.** – 82 males, 17 females.

**Distribution.** – Palaearctic Region (except for the extreme north) and eastern parts of North America.

**IUCN status.** – Least Concern.

**Xylota sylvarum** (Linnaeus, 1758)

**References.** – BECKER *et al.* (1910); DIRICKX (1994); SPEIGHT *et al.* (2018, 2020).

**Material examined.** – 5 males, 8 females.

**Distribution.** – Palaearctic Region.

**IUCN status.** – Least Concern.

## DISCUSSION

In the present study, we report 30 species from Corsica for the first time (including *Eumerus emarginatus*), as well as the genera *Anasimyia*, *Brachyopa*, *Lapposyrphus* Dušek & Láska, 1967, *Lejops* Rondani, 1857, *Melangyna* Verrall, 1901, *Neocnemodon*, and *Parhelophilus* Girschner, 1897. Among the studied hover flies, there are seven species that are endemic of Corsica (*Eumerus niehuisi*, *Eupeodes vandergooti*, *Riponnensis daccordii*) or have a distribution limited to the islands in the middle of the Mediterranean Sea, namely Corsica, Sardinia and Sicily (*Eumerus vandenberghaei*, *Merodon rubidiventris*, *Paragus ascoensis*, *P. sexarculus*).

Among the 131 sampled taxa, there are one Critically Endangered species, 10 Endangered species, six Vulnerable species, and five Near Threatened species. This means that near 13% of the hover fly species collected during the project *Our Planet Reviewed in Corsica 2019-2021* are threatened at the European level. We would also like to point out some remarkable records, such as the second record from Corsica of the Endangered *Doros destillatorius*, recently reported by CORNUEL-WILLERMOZ (2021), or the second record from Corsica of *Sericomyia silentis* since 1913 (KUNTZE, 1913). Furthermore, we report a new location for the Critically Endangered *Riponnensis daccordii* and corroborate that it is not extinct.

As mentioned by TOIROULT *et al.* (2023), the survey *Our Planet Reviewed in Corsica 2019-2021* did not sample evenly in Corsica. Thus, we can expect additional new species records for Corsica with more surveys on the island. We hope our results contribute to a better knowledge of the Syrphidae fauna of Corsica and help to make management decisions regarding these important pollinators.

### ORCID

Ximo Mengual :  <https://orcid.org/0000-0002-6185-9404>

Thomas Lebard :  <https://orcid.org/0000-0001-7538-9477>

Alexandre Cornuel-Willermoz :  <https://orcid.org/0009-0008-6158-3496>

**ACKNOWLEDGEMENTS.** – All of the material of this study was collected during the naturalist expedition *La Planète Revisitée (Our Planet Reviewed) en Corse 2019-2021*. This survey was organised by the Muséum national d'Histoire naturelle (MNHN, Paris) in collaboration with and funded by the Collectivité de Corse (CdC) and the Office Français de la Biodiversité (OFB). We are also grateful to our logistical partners who assisted with fieldwork in 2020: the Office de l'Environnement de la Corse (OCIC and CBNC), the Direction Régionale de l'Environnement de l'Aménagement et du Logement (DREAL) and the Conservatoire du Littoral (CdL).

We are indebted to the leaders of the expedition, Julien Touroult, François Dusoulier and Jean Ichter, for the thorough preparation and guidance. Thanks to the Diptera coordinator, Marc Pollet, for gathering and processing the pan trap and other Diptera samples, and providing us with the material used in this paper. We thank Frank van de Meutter for the identification of some *Cheilosia* specimens and two anonymous reviewers for the good suggestions and comments.

## REFERENCES

- AČANSKI J., VUJIĆ A., ŠAŠIĆ ZORIĆ L., RADENKOVIĆ S., DJAN M., MARKOV RISTIĆ Z. & STĀHLS G., 2022. – *Merodon chalybeus* subgroup: an additional piece of the *M. aureus* group (Diptera, Syrphidae) puzzle. *Annales Zoologici Fennici*, **59** : 79-109. <https://doi.org/10.5735/086.059.0109>
- ARACIL A., GRKOVIĆ A., PÉREZ-BAÑÓN C., KOČIŠ TUBIĆ N., JUAN A., RADENKOVIĆ S., VUJIĆ A. & ROJO S., 2023. – A new species of phytophagous flower fly (Diptera, Syrphidae), feeding on holoparasitic broomrape plants (Orobanchaceae) for the first time in Europe. *Arthropod-Plant Interactions*, **17** : 401-418. <https://doi.org/10.1007/s11829-023-09962-z>
- ARCAYA E., PÉREZ-BAÑÓN C., MENGUAL X., ZUBCOFF-VALLEJO J. J. & ROJO S., 2017. – Life table and predation rates of the syrphid fly *Allograpta exotica*, a control agent of the cowpea aphid *Aphis craccivora*. *Biological Control*, **115** : 74-84. <https://doi.org/10.1016/j.bioccontrol.2017.09.009>
- BARENDEGT A., ZEETERS T., VAN STEENIS W. & JONGEJANS E., 2022. – Forest hoverfly community collapse: abundance and species richness drop over four decades. *Insect Conservation and Diversity*, **15** : 510-521. <https://doi.org/10.1111/icad.12577>
- BARTSCH H., 2009. – Tvåvingar: Blomflugor - Diptera: Syrphidae: Eristalinae & Microdontinae. In : Nationalnyckeln till Sveriges flora och fauna. Uppsala : ArtDatabanken, DH 53b, 478 p.
- BECKER T., KUNTZE A., SCHNABL J. & VILLENEUVE E., 1910. – Dipterologische Sammelreise nach Korsika (Dipt.). Ausgeführt im Mai und Juni 1907 von Th. Becker, A. Kuntze, J. Schnabl und E. Villeneuve. *Deutsche Entomologische Zeitschrift*, **1910** : 635-665.
- BELLEFEUILLE Y., FOURNIER M. & LUCAS E., 2019. – Evaluation of two potential biological control agents against the Foxglove aphid at low temperatures. *Journal of Insect Science*, **19** : 2. <https://doi.org/10.1093/jisesa/iey130>
- BIGOT J. M. F., 1861. – Trois diptères nouveaux de la Corse. *Annales de la Société entomologique de France*, (4) **1** : 227-228.
- BIGOT J. M. F., 1862. – Diptères nouveaux de la Corse découverts dans la partie montagneuse de cette île par M. E. Bellier de la Chavignerie, pendant l'été de 1861. *Annales de la Société entomologique de France*, **2** : 109-114.
- BIRTELE D., 2011. – Contributo alla conoscenza dei Syrphidae della Sardegna (Diptera). *Conservazione habitat invertebrati*, **5** : 659-715.
- BOT S., HADRAVA J. & PENNARDS G., 2023. – The first confirmed records of *Riponnensis insignis* (Loew, 1843) (Diptera, Syrphidae) for Europe and an identification key to the European *Riponnensis* species. *Journaal van Syrphidae*, **2** (3) : 1-14. <https://doi.org/10.55710/1.ILFW2845>
- BOT S., MENGUAL X., VAN STEENIS J. & SKEVINGTON J. H., 2022. – A new species of the genus *Milesia* Latreille (Diptera: Syrphidae) from Crete. *European Journal of Taxonomy*, **846** : 110-125. <https://doi.org/10.5852/ejt.2022.846.1969>
- CHRONI A., GRKOVIĆ A., AČANSKI J., VUJIĆ A., RADENKOVIĆ S., VELIČKOVIĆ N., DJAN M. & PETANIDOU T., 2018. – Disentangling a cryptic species complex and defining new species within the *Eumerus minotauros* group (Diptera: Syrphidae), based on integrative taxonomy and Aegean palaeogeography. *Contributions to Zoology*, **87** : 197-225. <https://doi.org/10.1163/18759866-08704001>
- CLAUSSEN C. & TORP E., 1980. – Untersuchungen über vier europäische Arten der Gattung *Anasimyia* Schiner, 1864 (Insecta, Diptera, Syrphidae). *Mitteilungen aus dem Zoologischen Museum der Universität Kiel*, **1** (4) : 3-16, pl. I-IV.
- CLAUSSEN C., 1991. – Eine neue *Orthonevra* von Korsica (Diptera, Syrphidae). *Entomofauna*, **12** (16) : 205-212.

- CORNUEL-WILLERMOZ A., 2021. – Première mention de *Doros destillatorius* Mik, 1885 en Corse (Diptera Syrphidae). *L'Entomologiste*, **77** (5) : 289-292.
- CORNUEL-WILLERMOZ A., LEBARD T., BOT S. & MENGUAL X., 2023. – Découverte d'*Eumerus emarginatus* Loew, 1848 en Corse, une nouvelle espèce pour la faune de France (Diptera: Syrphidae). *L'Entomologiste*, **79** (4) : 245-250.
- DIRICKX H. G., 1994. – Atlas des Diptères syrphides de la région méditerranéenne. *Documents de Travail de l'Institut Royal des Sciences naturelles de Belgique*, **75** : 1-317.
- DOCZKAL D., 1996. – Description of two new species of the genus *Eumerus* (Diptera, Syrphidae) from Corsica. *Volucella*, **2** : 3-19.
- DOYLE T., HAWKES W. L. S., MASSY R., POWNEY G. D., MENZ M. H. M. & WOTTON K. R., 2020. – Pollination by hoverflies in the Anthropocene. *Proceedings of the Royal Society B: Biological Sciences*, **287** : 20200508. <https://doi.org/10.1098/rspb.2020.0508>
- DUNN L., LEQUERICA M., REID C. R. & LATTY, T., 2020. – Dual ecosystem services of syrphid flies (Diptera: Syrphidae): pollinators and biological control agents. *Pest Management Science*, **76** : 1973-1979. <https://doi.org/10.1002/ps.5807>
- DUŠEK J. & LÁSKA P., 1973. – Descriptions of five new European species of the genus *Metasyrphus* (Diptera: Syrphidae), with notes on variation within the species. *Acta entomologica bohemoslovaca*, **70** (6) : 415-426.
- FLEISCHMANN A., GONELLA P. M., ROJO S., & MENGUAL X., 2022. – Attracted to feed, not to be fed upon – on the biology of *Toxomerus basalis* (Walker, 1836), the kleptoparasitic ‘sundew flower fly’ (Diptera: Syrphidae). *Journal of Tropical Ecology*, **38** : 241-253. <https://doi.org/10.1017/S0266467422000128>
- FLEISCHMANN A., RIVADAVIA F., GONELLA P. M., PÉREZ-BAÑÓN C., MENGUAL X. & ROJO S., 2016. – Where is my food? Brazilian flower fly steals prey from carnivorous sundews in a newly discovered plant-animal interaction. *PLoS ONE*, **11** : e0153900. <https://doi.org/10.1371/journal.pone.0153900>
- GATTER W., EBENHÖH H., KIMA R., GATTER W. & SCHERER F., 2020. – 50-jährige Untersuchungen an migrierenden Schwebfliegen, Waffenfliegen und Schlupfwespen belegen extreme Rückgänge (Diptera: Syrphidae, Stratiomyidae; Hymenoptera: Ichneumonidae). *Entomologische Zeitschrift*, **130** : 131-142.
- GOELDLIN DE TIEFENAU P., 1976. – Révision du genre *Paragus* (Dipt. Syrphidae) de la région paléarctique occidentale. *Mitteilungen der schweizerischen Entomologischen Gesellschaft*, **49** : 79-108.
- GOELDLIN DE TIEFENAU P. & LUCAS J. A. W., 1981. – *Paragus* (Dipt., Syrphidae) de Corse et de Sardaigne. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft*, **54** : 389-397.
- GRKOVIĆ A., VUJIĆ A., CHRONI A., VAN STEENIS J., ĐAN M. & RADENKOVIĆ S., 2017. – Taxonomy and systematics of three species of the genus *Eumerus* Meigen, 1822 (Diptera: Syrphidae) new to southeastern Europe. *Zoologischer Anzeiger*, **270** : 176-192. <https://doi.org/10.1016/j.jcz.2017.10.007>
- GRKOVIĆ A., VUJIĆ A., RADENKOVIĆ S., CHRONI A. & PETANIDOU T., 2015. – Diversity of the genus *Eumerus* Meigen (Diptera, Syrphidae) on the eastern Mediterranean islands with description of three new species. *Annales de la Société entomologique de France* (N.S.), **51** (4) : 361-373. <https://doi.org/10.1080/00379271.2016.1144483>
- GROSSKOPF G., 2005. – Biology and life history of *Cheilosia urbana* (Meigen) and *Cheilosia psilophthalma* (Becker), two sympatric hoverflies approved for the biological control of hawkweeds (*Hieracium* spp.) in New Zealand. *Biological Control*, **35** : 142-154. <https://doi.org/10.1016/j.bicontrol.2005.06.013>
- HEO C. C., RAHIMI R., MENGUAL X., M. ISA M. S., ZAINAL S., KHOFAR P. N. & NAZNI W. A., 2020. – *Eristalinus arvorum* (Fabricius, 1787) (Diptera: Syrphidae) in human skull: a new fly species of forensic importance. *Journal of Forensic Sciences*, **65** : 276-282. <https://doi.org/10.1111/1556-4029.14128>
- HIPPA H., 1968. – Classification of the Palaearctic species of the genera *Xylota* Meigen and *Xylotomina* Shannon (Dipt., Syrphidae). *Annales entomologica Fennica*, **34** : 179-197.
- HURKMANS W., 1993. – A monograph of *Merodon* (Diptera: Syrphidae). Pt.1. *Tijdschrift voor Entomologie*, **136** : 147-234.
- INOUE D. W., LARSON B. M. H., SSYMANIK A. & KEVAN P. G., 2015. – Flies and Flowers III: ecology of foraging and pollination. *Journal of Pollination Ecology*, **16** : 115-133. [https://doi.org/10.26786/1920-7603\(2015\)15](https://doi.org/10.26786/1920-7603(2015)15)

- IUCN, 2023. – The IUCN Red List of Threatened Species. Version 2022-2. Available from <https://www.iucnredlist.org> [accessed 4.V.2023]
- JANKOVIĆ M. & RADENKOVIĆ S., 2021. – *Merodon aerarius*. The IUCN Red List of Threatened Species 2021: e.T149115186A149115188 [accessed 2.V.2023].  
<https://doi.org/10.2305/IUCN.UK.2021-3.RLTS.T149115186A149115188.en>
- KUNTZE A., 1913. – Dipterologische Sammelreise in Korsika des Herrn W. Schnuse in Dresden im Juni und Juli 1899. *Deutsche Entomologische Zeitschrift*, **1913** : 544-567.
- LAIR X., ROPARS L., SKEVINGTON J. H., KELSO S., GESLIN B., MINSSIEUX E. & NÈVE G., 2022. – Revision of the genus *Pelecocera* Meigen, 1822 (Diptera: Syrphidae) from France: taxonomy, ecology and distribution. *Zootaxa*, **5141** : 1-24. <https://doi.org/10.11646/zootaxa.5141.1.1>
- LARDÉ G., 1989. – Investigation on some factors affecting larval growth in a coffee-pulp bed. *Biological Wastes*, **30** : 11-19. [https://doi.org/10.1016/0269-7483\(89\)90139-0](https://doi.org/10.1016/0269-7483(89)90139-0)
- LARDÉ G., 1990. – Growth of *Ornidia obesa* (Diptera: Syrphidae) larvae on decomposing coffee pulp. *Biological Wastes*, **34** : 73-76. [https://doi.org/10.1016/0269-7483\(90\)90144-H](https://doi.org/10.1016/0269-7483(90)90144-H)
- LEBARD T., CANUT M. & SPEIGHT M. C. D., 2019. – Première observation en France d'*Ischiodon aegyptius* (Wiedemann, 1830) et découverte en Corse d'*Eumerus narcissi* Smith, 1928 (Diptera, Syrphidae). *Revue Française d'Entomologie Générale*, **1** (3) : 203-210.
- MAGNI P., PÉREZ-BAÑÓN C., BORRINI M. & DADOUR I., 2013. – *Syritta pipiens* (Diptera: Syrphidae), a new species associated with human cadavers. *Forensic Science International*, **231** : e19-e23.  
<https://doi.org/10.1016/j.forsciint.2013.05.023>
- MARCOS-GARCÍA M. Á., 1986. – Citas de interés del género *Paragus* Latreille, 1804 en la Península Ibérica y descripción de *Paragus vandergooti* sp. nov. (Diptera, Syrphidae) (p. 765-772). *Actas de las VIII Jornadas de la Asociación española de Entomología, Sevilla*.
- MARCOS-GARCÍA M. Á., MAZÁNEK L., LÁSKA P., BIČÍK V. & ROJO S., 2000. – Description of the male of *Eupeodes lucasi* (Marcos-García & Láska, 1983) and biological data on the species (Diptera, Syrphidae). *Volucella*, **5** : 129-138.
- MARCOS-GARCÍA M. Á. & ROJO S., 1994. – *Paragus hyalopteri* n. sp. an aphidophagous hoverfly (Dipt.: Syrphidae) attacking the mealy plum aphid (Hom.: Aphidae). *Entomophaga*, **39** (1) : 99-106.
- MARCOS-GARCÍA M. Á., VUJIĆ A. & MENGUAL X., 2007. – Revision of Iberian species of the genus *Merodon* (Diptera: Syrphidae). *European Journal of Entomology*, **104** : 531-572.  
<https://doi.org/10.14411/eje.2007.073>
- MAZÁNEK L., RICARTE SABATER A. R., VUJIĆ A., MILIĆ M. & NEDELJKOVIĆ Z., 2021. – *Eupeodes vandergooti*. The IUCN Red List of Threatened Species 2021: e.T149171568A149171571 [accessed 02 May 2023]. <https://doi.org/10.2305/IUCN.UK.2021-3.RLTS.T149171568A149171571.en>
- MENGUAL X., BOT S., CHKHARTISHVILI T., REIMANN A., THORMANN J. & VON DER MARK L., 2020. – Checklist of hover flies (Diptera, Syrphidae) of the Republic of Georgia. *ZooKeys*, **916** : 1-123.  
<https://doi.org/10.3897/zookeys.916.47824>
- MENGUAL X., KAZERANI F., ASGHAR TALEBI A. & GILASIAN E., 2015a. – A revision of the genus *Pelecocera* Meigen with the description of the male of *Pelecocera persiana* Kuznetsov from Iran (Diptera: Syrphidae). *Zootaxa*, **3947** : 99-108. <https://doi.org/10.11646/zootaxa.3947.1.6>
- MENGUAL X., MAYER C., BURT T. O., MORAN K. M., DIETZ L., NOTTEBROCK G., PAULI T., YOUNG A. D., BRASSEUR M. V., KUKOWKA S., KELSO S., ETZBAUER C., BOT S., HAUSER M., JORDAENS K., MIRANDA G. F. G., STÄHLS G., VAN STEENIS W., PETERS R. S. & SKEVINGTON J. H., 2023. – Systematics and evolution of predatory flower flies (Diptera: Syrphidae) based on exon-capture sequencing. *Systematic Entomology*, **48** (2) : 250-277. <https://doi.org/10.1111/syen.12573>
- MENGUAL X., STÄHLS G. & ROJO S., 2015b. – Phylogenetic relationships and taxonomic ranking of pipizine flower flies (Diptera: Syrphidae) with implications for the evolution of aphidophagy. *Cladistics*, **31** : 491-508. <https://doi.org/10.1111/cla.12105>
- MILIĆ M. & GRKOVIĆ A., 2021. – *Eumerus niehuisi*. The IUCN Red List of Threatened Species 2021: e.T149170245A149170247 [accessed 02 May 2023].  
<https://doi.org/10.2305/IUCN.UK.2021-3.RLTS.T149170245A149170247.en>

- MOERKENS R., BOONEN S., WÄCKERS F. L. & PEKAS A., 2021. – Aphidophagous hoverflies reduce foxglove aphid infestations and improve seed set and fruit yield in sweet pepper. *Pest Management Science*, **77** (6) : 2690-2696. <https://doi.org/10.1002/ps.6342>
- MORALES G. E. & WOLFF M., 2010. – Insects associated with the composting process of solid urban waste separated at the source. *Revista Brasileira de Entomologia*, **54** (4) : 645-653. <https://doi.org/10.1590/S0085-56262010000400017>
- MORAN K. M., SKEVINGTON J. H., KELSO S., MENGUAL X., JORDAENS K., YOUNG A. D., STÅHLS G., MUTIN V., BOT S., VAN ZUIJEN M., ICHIGE K., VAN STEENIS J., HAUSER M. & VAN STEENIS W., 2022. – A multigene phylogeny of the eristaline flower flies (Diptera: Syrphidae), with emphasis on the subtribe Criorhinina. *Zoological Journal of the Linnean Society*, **194** : 120-135. <https://doi.org/10.1093/zoolinnean/zlab006>
- PAULI T., BURT T., MEUSEMANN K., BAYLESS K., DONATH A., PODSIADLOWSKI L., MAYER C., KOZLOV A., VASILIKOPOULOS A., LIU S., ZHOU X., YEATES D., MISOF B., PETERS R. S. & MENGUAL X., 2018. – New data, same story: phylogenomics does not support Syrphoidea (Diptera: Syrphidae, Pipunculidae). *Systematic Entomology*, **43** (3) : 447-459. <https://doi.org/10.1111/syen.12283>
- PELLMANN, H., 1998. – Die Gattung *Brachyopa* Meigen, 1822 (Insecta, Diptera, Syrphidae) in entomologischen Sammlungen sächsischer Museen und die Möglichkeit der Artunterscheidung anhand der Genitalien der Männchen. *Studia dipterologica*, **5** (1) : 95-112.
- PÉREZ-BAÑÓN C., ROJAS C., VARGAS M., MENGUAL X. & ROJO, S., 2020. – A world review of reported myiasis caused by flower flies (Diptera: Syrphidae), including the first case of human myiasis from *Palpada scutellaris* (Fabricius, 1805). *Parasitology Research*, **119** : 815–840. <https://doi.org/10.1007/s00436-020-06616-4>
- PÉREZ-LACHAUD G., JERVIS M. A., REEMER M. & LACHAUD, J.-P., 2014. – An unusual, but not unexpected, evolutionary step taken by syrphid flies: the first record of true primary parasitoidism of ants by Microdontinae. *Biological Journal of the Linnean Society*, **111** : 462-472. <https://doi.org/10.1111/bij.12220>
- POPOVIĆ D., AČANSKI J., DJAN M., OBREHT D., VUJIĆ A. & RADENKOVIĆ S., 2015. – Sibling species delimitation and nomenclature of the *Merodon avidus* complex (Diptera: Syrphidae). *European Journal of Entomology*, **112** (4) : 790-809. <https://doi.org/10.14411/eje.2015.100>
- POTTS S., DAUBER J., HOCHKIRCH A., OTEMAN B., ROY D., AHNRE K., BIESMEIJER K., BREEZE T., CARVELL C., FERREIRA C., FITZPATRICK Ú., ISAAC N. J. B., KUSSAARI M., LJUBOMIROV T., MAES J., NGO H., PARDO A., POLCE C., QUARANTA M., SETTELE J., SORG M., STEFANESCU C. & VUJIĆ A., 2021. – *Proposal for an EU Pollinator Monitoring Scheme. EUR 30416 EN, JRC122225*. Luxembourg : Publications Office of the European Union, 310 p. <https://doi.org/10.2760/881843>
- REEMER M., HAUSER M. & SPEIGHT M. C. D., 2004. – The genus *Myolepta* Newman in the West-Palaearctic region (Diptera, Syrphidae). *Studia dipterologica*, **11** (2) : 553-580.
- RICARTE A., NEDELJKOVIĆ Z., ROTHERAY G. E., LYSZKOWSKI R. M., HANCOCK E. G., WATT K., HEWITT S. M., HORSFIELD D. & WILKINSON G., 2012. – Syrphidae (Diptera) from the Greek island of Lesvos, with description of two new species. *Zootaxa*, **3175** : 1-23. <https://doi.org/10.11646/zootaxa.3175.1.1>
- RICARTE A., SOUBA-DOLS G. J., HAUSER M. & MARCOS-GARCÍA M.-Á., 2017. – A review of the early stages and host plants of the genera *Eumerus* and *Merodon* (Diptera: Syrphidae), with new data on four species. *Plos ONE*, **12** (12): e0189852. <https://doi.org/10.1371/journal.pone.0189852>
- RIZZA A., CAMPOBASSO G., DUNN P. H. & STAZI, M., 1988. – *Cheilosia corydon* (Diptera: Syrphidae), a candidate for the biological control of musk thistle in North America. *Annals of the Entomological Society of America*, **81** : 225-232. <https://doi.org/10.1093/aesa/81.2.225>
- ROTHERAY G. E., 1993. – *Colour Guide to Hoverfly Larvae (Diptera, Syrphidae) in Britain and Europe. Dipterists Digest No. 9*. Sheffield : Derek Whiteley, 156 p.
- ROTHERAY G. E. & GILBERT, F., 2011. – *The natural history of hoverflies*. Ceredigion : Forrest Text, 334 p.
- SARTHOU J.-P., DEHOND F., DUSSAIX C., LIBERT P.-N., VANAPPELGHEM C. & VAN DE WEYER G., 2004. – Contribution à la connaissance des Syrphes de 27 départements français ; deux espèces nouvelles pour la France (Diptera, Syrphidae et Microdonthidae). *Bulletin de la Société entomologique de France*, **109** (2) : 169-174. <https://doi.org/10.3406/bsef.2004.16102>

- SCHINER J. R., 1857. – Diptera Austriaca. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien*, 7 : 279-506.
- SHEPPARD A. W., AESCHLIMANN J.-P., SAGLIOCCO J.-L. & VITOU, J., 1995. – Below-ground herbivory in *Carduus nutans* (Asteraceae) and the potential for biological control. *Biocontrol Science and Technology*, 5 (3) : 261-270. <https://doi.org/10.1080/09583159550039729>
- SHORTHOUSE D. P., 2010. – SimpleMappr, an online tool to produce publication-quality point maps. Available from <https://www.simplemappr.net> [accessed 10.X.2023]
- SKEVINGTON J. H., LOCKE M. M., YOUNG A. D., MORAN K., CRINS W. J. & MARSHALL, S. A., 2019. – *Field Guide to the Flower Flies of Northeastern North America*. Princeton : Princeton University Press, 511 p. <https://doi.org/10.2307/j.ctv7xbrvz>
- SMIT J., 2014. – Two new species of the genus *Callicera* Panzer (Diptera: Syrphidae) from the Palaearctic Region. *Zootaxa*, 3779 (5) : 585-590. <https://doi.org/10.11646/zootaxa.3779.5.8>
- SOMMAGGIO D., 2001. – The species of the genus *Chrysotoxum* Meigen, 1822 (Diptera: Syrphidae) described by Giglio Tos. *Bollettino del Museo Regionale di Scienze Naturali di Torino*, 18 : 115-126.
- SOMMAGGIO D., 2002. – *Paragus gorgus* Vujić & Radenković, 1999: a junior synonym of *P. majoranae* Rondani, 1857, and reinstatement of *P. pecchiolii* Rondani, 1857 (Diptera, Syrphidae). *Volucella*, 6 : 53-56.
- SPEIGHT M. C. D., 1988. – *Doros destillatorius*, espèce nouvelle pour la France, avec désignation des types des deux espèces européennes du genre *Doros*, description de leurs pupes et clés de détermination des adultes et des pupes. *Bulletin de la Société entomologique de France*, 92 : 193-200. <https://doi.org/10.3406/bsef.1987.17496>
- SPEIGHT M. C. D., 1991. – *Callicera aenea*, *C. aurata*, *C. fagesii* and *C. macquartii* redefined, with a key to and notes on the European *Callicera* species (Diptera: Syrphidae). *Dipterists Digest*, 10 : 1-25.
- SPEIGHT M. C. D., 2014. – *Species accounts of European Syrphidae (Diptera), 2014. Syrph the Net, the database of European Syrphidae (Diptera)*, Vol. 78. Dublin : Syrph the Net publications, 321 p.
- SPEIGHT M. C. D., 2018. – *Species accounts of European Syrphidae, 2018. Syrph the Net, the database of European Syrphidae (Diptera)*, Vol. 103. Dublin : Syrph the Net publications, 302 p.
- SPEIGHT M. C. D., 2020a. – *StN key for the identification of the genera of European Syrphidae (Diptera). Syrph the Net, the database of European Syrphidae (Diptera)*, Vol. 105. Dublin : Syrph the Net publications, 46 p.
- SPEIGHT M. C. D., 2020b. – *Species accounts of European Syrphidae, 2020. Syrph the Net, the database of European Syrphidae (Diptera)*, Vol. 104. Dublin : Syrph the Net publications, 314 p.
- SPEIGHT M. C. D., CASTELLA E. & SARTHOU J.-P., 2020. – StN 2020 – In : Speight, M. C. D., Castella, E., Sarthou, J.-P. & Vanappelghem, C. (eds), *Syrph the Net on CD, Issue 12*. ISSN 1649-1917. Dublin : Syrph the Net publications, [excel spreadsheet].
- SPEIGHT M. C. D., CLAUSSSEN C. & HURKMAN W., 1998. – Révision des syrphes de la faune de France : III - Liste alphabétique des espèces des genres *Cheilosia*, *Eumerus* et *Merodon* et Supplément (Diptera, Syrphidae). *Bulletin de la Société entomologique de France*, 103 : 403-414. <https://doi.org/10.3406/bsef.1998.17451>
- SPEIGHT M. C. D., FISLER L., PÉTREMAND G. & HAUSER M., 2021. – *A key to the males of the Eumerus species known from Switzerland & surrounding parts of Central Europe (Diptera: Syrphidae). Syrph the Net, the database of European Syrphidae (Diptera)*, Vol. 112. Dublin : Syrph the Net publications, 36 p.
- SPEIGHT M. C. D. & LANGLOIS D., 2020. – *Clés des mâles des espèces françaises de Merodon, 2020 (Diptera: Syrphidae). Syrph the Net, the database of European Syrphidae (Diptera)*, Vol. 110. Dublin : Syrph the Net publications, 60 p.
- SPEIGHT M. C. D. & LEBARD T., 2020 – Données de syrphes nouvelles pour les départements français. *Bulletin de la Société Linnéenne de Bordeaux*, 155 : 341-354.
- SPEIGHT M. C. D. & LEBARD T., 2022. – Quelques additions à la liste des syrphes connus dans le département du Gard, avec une mise à jour de la clef des taxons du groupe *Chrysotoxum intermedium* en France (Diptera : Syrphidae). *Revue Française d'Entomologie Générale*, 4 : 15-31.

- SPEIGHT M. C. D. & SARTHOU J.-P., 2006. – Révision de la liste des Diptères Syrphidae et Microdontidae de France métropolitaine et de Corse : 505 espèces confirmées dont 13 nouvelles pour cette faune. *Bulletin de la Société entomologique de France*, **111** (1) : 11-20. <https://doi.org/10.3406/bsef.2006.16277>
- SPEIGHT M. C. D. & SARTHOU J.-P., 2017. – StN keys for the identification of the European species of various genera of Syrphidae. *Syrph the Net, the database of European Syrphidae (Diptera)*, Vol. 99. Dublin : Syrph the Net publications, 139 p.
- SPEIGHT M. C. D., SARTHOU J.-P., VANAPPELGHEM C. & SARTHOU V., 2018. – Maps of the departmental distribution of syrphid species in France / Cartes de distribution départemental des syrphes de France (Diptera: Syrphidae). *Syrph the Net, the database of European Syrphidae (Diptera)*, Vol. 100. Dublin : Syrph the Net publications, 80 p.
- SSYMANIK A. & KEARNS C., 2009. – Flies-pollinators on two wings (p. 39-52). In : Ssymank A., Hamm A. & Vischer-Leopold M. (eds), *Caring for pollinators – safeguarding agro-biodiversity and wild plant diversity*. Bonn : Bundesamt für Naturschutz.
- STÅHLS G., 2022. – Taxo-Fly - an EU-funded project gathering taxonomic information for all European hoverfly species. Available from <https://www.luomus.fi/en/taxo-fly> [accessed 18.IV.2023]
- TOUROULT J., ICHTER J., POLLET M., PASCAL O., POIRIER E., ROUGERIE R., DECHERF B., ANDREI-RUIZ M.-C., HUGOT L. & DUSOULIER F., 2023. – Our Planet Reviewed in Corsica 2019-2021: a large-scale survey of neglected biodiversity on a Mediterranean island. *Bulletin de la Société entomologique de France*, **128** (4) : 353-382. [https://doi.org/10.32475/bsef\\_2285](https://doi.org/10.32475/bsef_2285)
- VAN DER GOOT V. S., 1961. – Zweefvliegenvangst op Corsica. *Entomologische Berichten*, **21** : 219-223.
- VAN DER GOOT V. S., 1964. – Summer records of syrphidae (diptera) from Sicily, with field notes and descriptions of new species. *Zoologische Mededelingen*, **39** (42) : 414-432.
- VAN DER GOOT V. S., 1968. – A new *Eumerus* (Syrphidae, Diptera) species from Corsica. *Entomologische Berichten*, **28** : 219-220.
- VAN ECK A. & MENGUAL X., 2021. – Review of the genus *Pelecocera* Meigen, 1822 (Diptera, Syrphidae) in the Palaearctic with the description of a new species from Cyprus. *Beiträge zur Entomologie*, **71** (2) : 321-343. <https://doi.org/10.21248/contrib.entomol.71.2.321-343>
- VAN STEENIS J., RICARTE A., VUJIĆ A., BIRTELE D., & SPEIGHT M. C. D., 2016. – Revision of the West-Palaearctic species of the tribe Cerioidini (Diptera, Syrphidae). *Zootaxa*, **4196** (2) : 151-209. <https://doi.org/10.11164/zootaxa.4196.2.1>
- VAN STEENIS W., VAN STEENIS J. & VAN DER ENT L.-J., 2021. – *Riponnensis daccordii*. The IUCN Red List of Threatened Species 2021: e.T149165101A149165112 [accessed 3.V.2023]. <https://doi.org/10.2305/IUCN.UK.2021-3.RLTS.T149165101A149165112.en>
- VAN VEEN M. P., 2010. – *Hoverflies of Northwest Europe: identification keys to the Syrphidae. 2nd edition*. Utrecht : KNNV Publishing, 248 p.
- VILLENEUVE J., 1909. – Description d'un nouveau Syrphide. *Wiener Entomologische Zeitung*, **28** : 338-339.
- VILLENEUVE J., 1912. – Notes sysnonymiques. *Wiener Entomologische Zeitung*, **31** : 96-97.
- VUJIĆ A., GILBERT F., FLINN G., ENGLEFIELD E., VARGA Z., FERREIRA C. C., EGGERT F., WOOLCOCK S., BÖHM M., VBRA J., MERGY R., SSYMANIK A., VAN STEENIS W., ARACIL A., FÖLDÉSI R., GRKOVIĆ A., MAZANEK L., NEDELJKOVIĆ Z., PENNARDS G. W. A., PÉREZ C., RADENKOVIĆ S., RICARTE A., ROJO S., STÅHLS G., VAN DER ENT L.-J., VAN STEENIS J., BARKALOV A., CAMPOY A., JANKOVIĆ M., LIKOV L., LILLO I., MENGUAL X., MILIĆ D., MILIĆIĆ M., NIELSEN T., POPOV G., ROMIG T., ŠEBIĆ A., SPEIGHT M., TOT T., VAN ECK A., VESELIĆ S., ANDRIC A., BOWLES P., DE GROOT M., MARCOS-GARCÍA M. Á., HADRAVA J., LAIR X., MALIDŽAN S., NIÈVE G., OBREHT VIDAKOVIC D., POPOV S., SMIT J. T., VAN DE MEUTTER F. & VELIČKOVIĆ N., 2022. – *The European Red List of Hoverflies*. Brussels : European Commission, viii + 96 p.
- VUJIĆ A., NEDELJKOVIĆ Z., HAYAT R., DEMİRÖZER O., MENGUAL X. & KAZERANI F., 2017. – New data on the genus *Chrysotoxum* Meigen (Diptera: Syrphidae) from North-East Turkey, Armenia, Azerbaijan and Iran including descriptions of three new species. *Zoology in the Middle East*, **63** : 250-268. <https://doi.org/10.1080/09397140.2017.1349241>
- VUJIĆ A., RADENKOVIĆ S., LIKOV L. & VESELIĆ S., 2021. – Taxonomic complexity in the genus *Merodon* Meigen, 1803 (Diptera, Syrphidae). *ZooKeys*, **1031** : 85-124. <https://doi.org/10.3897/zookeys.1031.62125>

- VUJIĆ A., RADENKOVIĆ S., LIKOV L., ANDRIĆ A., JANKOVIĆ M., AČANSKI J., POPOV G., DE COURCY WILLIAMS M., ŠAŠIĆ ZORIĆ L. & DJAN M., 2020b. – Conflict and congruence between morphological and molecular data: revision of the *Merodon constans* group (Diptera: Syrphidae). *Invertebrate Systematics*, **34** (4) : 406-448. <https://doi.org/10.1071/IS19047>
- VUJIĆ A., SPEIGHT M., DE COURCY WILLIAMS M. E., ROJO S., STĀHLS G., RADENKOVIĆ S., LIKOV L., MILIĆIĆ M., PÉREZ-BAÑÓN C., FALK S. & PETANIDOU T., 2020a. – *Atlas of the Hoverflies of Greece*. Leiden : Brill, 384 p. <https://doi.org/10.1163/9789004334670>
- VUJIĆ A., STĀHLS, G., AČANSKI, J., BARTSCH, H., BYGEBJERG, R. & STEFANOVIĆ, A., 2013. – Systematics of Pipizini and taxonomy of European *Pipiza* Fallén: molecular and morphological evidence (Diptera, Syrphidae). *Zoologica Scripta*, **42** : 288-305. <https://doi.org/10.1111/zsc.12005>
- YOUNG A. D., LEMMON A. R., SKEVINGTON J. H., MENGUAL X., STĀHLS G., REEMER M., JORDAENS K., KELSO S., LEMMON E. M., HAUSER M., DE MEYER M., MISOF B. & WIEGMANN B. M., 2016. – Anchored enrichment dataset for true flies (order Diptera) reveals insights into the phylogeny of flower flies (family Syrphidae). *BMC Evolutionary Biology*, **16** : 143. <https://doi.org/10.1186/s12862-016-0714-0>
-